

Mixture of natural antioxidants can improve *Spodoptera littoralis* (Boisd.) nucleopolyhedrovirus, (Baculoviridae) stability under Egyptian field conditions

El-Helaly, A.

Dept. of Econ. Entomol. and Pesticides, Fac. Agric., Cairo Univ., Giza, Egypt



ABSTRACT

Four promising natural plants previously proved to prolong the activity of *Spodoptera littoralis* (Boisd.) nucleopolyhedrovirus (*SpliNPV*) suspension under Egyptian field conditions, examined alone or in combination with virus in six mixtures. Screening results were based on bioassay. Different parameters were investigated to prove the value of protection, % of mortality, reduction %, LIT₅₀, Potency and Original Activity Remaining (OAR). DPPH assay was investigated for all treatments. The obtained results showed that, virus mixed with cacao + green coffee treatment gave the highest LIT₅₀ (203.474 hours) while scored only (20.172 hours) with virus alone treatment. Original activity remaining indicated that *SpliNPV* mixed with both cacao + green coffee treatment preserved 100% of its activity for five days while it gave only 1.04 % for *SpliNPV* alone treatment and gave 41.06, 31.17, 17.65 and 42.56 with cacao, green coffee, moringa and both cacao + moringa additives; respectively five days post investigation. This investigation suggests that mixing natural derived antioxidants provides good protection to baculovirus at Egyptian sunny conditions.

Keywords: antioxidants, DPPH, OAR, nucleopolyhedrovirus, ultraviolet, *Spodoptera littoralis* NPV.

INTRODUCTION

The Biological control method is an important component of integrated pest management IPM. Baculoviruses biopesticides have many advantages as tool in IPM, (Ravensberg, 2011). However, baculoviruses, like others biopesticides, present some difficulties, such as short field stability (Mills and Kean, 2010; Regnault-Roger, 2012). More than 60 baculovirus bio-pesticides have been used to control diverse insect pest all around the world (Moscardi, 1999). The treated areas increased up to more than two million hectares in 2010-2012 (Yang *et al.*, 2012). Antioxidants derived from natural plants proved to elongate virus stability last decade. The first record of the use of plant extracts to increase the persistence of insect viruses was by Shapiro *et al.* (2007a, b). Both of the green tea and black tea were reported to be UV protective additive to the beet armyworm nucleopolyhedrovirus (Shapiro *et al.*, 2008 and El Salamouny *et al.*, 2009) Mango leaf extract (Deotale *et al.*, 2007) Cacao, green coffee, green and red cabbage (El-Helaly *et al.*, 2009 and El-Helaly *et al.*, 2013) Moringa and Rice bran (El-Helaly, 2013).

The present study was aimed to evaluate in both laboratory and field tests the role of addition the mixtures of the most promising previous additives Moringa, green tea, green coffee cacao and used *SpliMNPV* alone against *S. littoralis*.

MATERIALS AND METHODS

Test Insect

The Egyptian cotton leaf worm, *Spodoptera littoralis* (Boisd.) established on the semi-synthetic diet described by Shorey and Hale (1965) used in the present investigation.

Virus inocula

Spodoptera littoralis (Boisd.) multiple embedded nucleopolyhedrovirus (*SpliMNPV*), Egyptian isolate.

UV-Protective additives

Four different plant-derived materials (Moringa, green tea, green coffee and cacao) were evaluated as

UV-protective additives to *SpliMNPV* suspension. Alone or in mixtures as follows (cacao + moringa, cacao + green coffee, cacao + green tea, moringa + green coffee, moringa + green tea and green coffee + green tea) One gram of each dry plant material was soaked in 100 ml distilled water for 24 hours then blended and filtrated through three layers of muslin and the filtrate was the stock additive to the tested virus concentration (1×10^9 PIB's/ml) to prepare a final concentration of 1% of the material additive according to the method described by Shapiro *et al.*, (2008) for additives alone, for mixtures half gram of each additive was soaked.

Simulated UV radiation in the sunlight and bioassay

Sunlight UV was simulated using UV lamps according to (Huber and Ludcke, 1996). 50 μ l was spread inside a Petri dish (10 cm in diameter), virus film were exposed to the tested irradiation source, after exposures of virus treatment to UV irradiation. The polyhedra deposits in the Petri-dish were resuspended in 10 ml distilled water for use in bioassay tests. Two mls of collected PIB's suspension were applied on the surface of 50 ml semi-artificial diet. An un-treated control treated, with only distilled water. Neonate test larvae, of each treatment, were allowed to feed on the treated diet surface till pupation. Mortality recorded up to 14 days post application (Fritsch and Huber, 1985). Treated insects were laboratory maintained at 25 ± 2 °C and 65 ± 5 R. H.

Field experiment

The promising additives resulted from laboratory tests were further evaluated under outdoors field conditions. For this purpose, about 1/3 feddan of tomatoes (1 feddan = 4200 m²) was used and one small scale field test was set up to confirm laboratory results. The virus inocula *SpliNPV* was adjusted to 10^8 OBs/ml (=LC₉₀₋₉₅). At the time of field application, the virus and tested additives were mixed together and transferred into a hand sprayer. Virus suspension treatments were applied separately to tomato foliage using half liter hand sprayer. Untreated leaves and virus treated leaves were randomly collected at 0, 1, 2, 4, 7, and 10 days post application and kept individually in plastic bags at room

temperature until tested. Each leaf was placed into a glass bottle, on which 10 neonate larvae were allowed to feed for 24 hr. before transferred daily to fresh leaves from the same treatment. Larval mortality was recorded daily until day 14. (Shapiro *et al.*, 2008)

DPPH assay

The measurement of the DPPH radical scavenging activity was performed according to Brand-Williams *et al.*, 1995. The scavenging activity percentage was determined according to Mensor *et al.*, 2001.

Statistical analysis

Concentration-mortality regressions were calculated to determine the effectiveness of tested material as UV protective additives for the *SpliMNPV*. Slope and LC_{50s} values were calculated according to the method described by Finney (1971). Original activity remaining percentages (OARs %) were determined for each treatment according to Muro and Paul, (1985) in which NPV- caused larval mortality post UV exposure were divided by NPV- caused larval mortality pre UV exposure and multiplied by 100. for DPPH The experiment was done in triplicate for each substance.

The results were expressed as percentage decrease with respect to control values and compared by one-way ANOVA and Turkey's test. A difference was considered statistically significant if p<0.05.

RESULTS

Screening results revealed loosing of more than 80 % of virus alone treatment activity five hours post application, where it gave 48.00, 33.33, 25.00, 14.00 mortality % after 30, 60, 180, 300; respectively. (Table 1, Fig 1) while other treatments preserved the virus activity *SpliNPV* mixed with cacao, green tea, green coffee and moringa treatments gave 49.00, 28.12, 39.39, 40.20 mortality %; respectively. mixing natural antioxidants increase the mortality percentage when *SpliNPV* was mixed with cacao + green coffee, cacao + moringa, moringa+ green coffee and moringa+ green tea where it increased the mortality % to 62.10, 55.67, 45.83 and 40.00; respectively, mixing *SpliNPV* with cacao + green tea or with green coffee + green tea gave 30.20 or 23.65 respectively.

Table (1) Average rates of mortality among *S littoralis* neonate larvae treated with *SpliNPV* either alone or in combination with different antioxidants or their mixture, all exposed to different UV irradiation periods.

Irradiation exposure period (hours)	Mortality % among larvae tested with <i>SpliNPV</i> alone or mixed the indicated additives at 1%											
	<i>SpliNPV</i> alone	cacao	Green tea	green coffee	moringa	cacao+ moringa	cacao + green coffee	cacao+ green tea	Moringa + green coffee	moringa+ green tea	Green coffee + green tea	
Zero time	99.00 (99/100)	93.93 (93/99)	98.96 (96/97)	93.87 (92/98)	98.00 (98/100)	93.93 (93/99)	99.00 (99/100)	100.00 (98/98)	92.00 (92/100)	93.00 (93/100)	96.00 (96/100)	
0.5	48.00 (48/100)	93.75 (90/96)	98.96 (71/100)	72.44 (71/98)	85.41 (82/96)	93.00 (93/100)	100.00 (98/98)	71.00 (71/100)	89.69 (87/97)	76.00 (76/100)	69.00 (69/100)	
1	33.33 (32/96)	80.00 (80/100)	61.85 (60/97)	69.00 (69/100)	72.16 (70/97)	90.00 (90/100)	97.00 (97/100)	66.00 (66/100)	79.38 (77/97)	70.00 (70/100)	58.76 (57/97)	
3	25.00 (25/100)	71.00 (71/100)	39.00 (39/100)	43.00 (43/100)	64.58 (62/96)	76.04 (73/96)	82.29 (79/96)	31.00 (31/100)	68.00 (68/100)	61.85 (60/97)	39.00 (39/100)	
5	14.00 (14/100)	49.00 (49/100)	28.12 (27/96)	39.39 (39/99)	40.20 (39/97)	55.67 (54/97)	62.10 (59/95)	30.20 (29/96)	45.83 (44/96)	40.00 (40/100)	23.65 (22/93)	
Control*	0.00 (0/100)	0.00 (0/100)	0.00 (0/99)	0.00 (0/99)	0.00 (0/100)	1.03 (1/97)	0.00 (0/97)	0.00 (0/96)	0.00 (0/95)	0.00 (0/96)	0.00 (0/100)	
LIT ₅₀	70.561	411.241	321.455	412.122	390.247	496.231	851.231	439.211	532.177	536.211	211.11	




Fig (1) Average rates of mortality among *S littoralis* neonate larvae treated with *SpliNPV* either alone or in combination with different antioxidants or their mixtures, all exposed to 300 min UV irradiation periods.

The obtained results represented in (Table 2 and Fig 2) showed that virus alone treatment gave the highest rates in reduction where it gave 85.00% while it gave 44.93, 70.85, 54.48, 57.8, 38.26, 36.9, 69.8, 46.17, 53.00 and 72.35 % with cacao, green tea, green coffee, moringa, both of cacao + moringa, cacao + green coffee, cacao + green tea, moringa + green coffee, moringa +

green coffee and green coffee + green tea; respectively 300 min post application. The highest LIT₅₀ found when that virus mixed with cacao + green coffee, where its activity lasts for 851.231 min (14.18 Hours) while it lasts for only 70.561 min (1.17 hours) with virus alone treatment.

Table (2) Average rates of reduction among *S littoralis* neonate larvae treated with *SpliNPV* either alone or in combination with different antioxidants or their mixture, all exposed to different UV irradiation periods.

Irradiation exposure period (hours)	Mortality % among larvae tested with <i>SpliNPV</i> alone or mixed the indicated additives										
	<i>SpliNPV</i> alone	cacao	green tea	green coffee	moringa	cacao+ moringa	cacao + green coffee	cacao+ green tea	moringa+ green coffee	moringa + green tea	green coffee + green tea
Zero time	99.00	93.93	98.96	93.87	98.00	93.93	99.00	100.00	92.00	93.00	96.00
0.5	51.00	0.18	0.00	21.43	12.59	0.93	-----	29.00	2.31	17.00	27.00
1	65.67	13.93	37.11	24.87	28.84	3.93	2.00	34.00	12.62	23.00	37.24
3	74.00	22.93	59.96	50.87	33.42	17.89	16.71	69.00	24.00	31.15	57.00
5	85.00	44.93	70.84	54.48	57.8	38.26	36.9	69.8	46.17	53.00	72.35




Fig (2) Average rates of reduction among *S littoralis* neonate larvae treated with *SpliNPV* either alone or in combination with different antioxidants or their mixture, all exposed to different UV irradiation periods.

Original activity remaining for *SpliNPV* alone treatment gave 48.48, 33.66, 25.25, 14.14 % half hour, one, three and five hours post application; respectively, while it increased to 99.80, 85.16, 75.58 and 52.16 % when *SpliNPV* mixed with cacao and it gave 100.00, 62.50, 39.40 and 28.41 when *SpliNPV* mixed with green tea, 77.17, 73.50, 45.80 and 41.96 % when *SpliNPV* mixed with green coffee, 87.15, 73.63, 65.89 and 41.02 % when *SpliNPV* mixed with moringa, 99.00, 95.81, 80.95 and 59.26 % when *SpliNPV* mixed with cacao+

moringa, 100.00, 97.97, 83.12 and 62.72 % when *SpliNPV* mixed with cacao + green coffee, 71.00, 66.00, 31.00 and 30.20 % when *SpliNPV* mixed with cacao + green tea, 97.78, 86.28, 73.91 and 49.81 % when *SpliNPV* mixed with moringa + green coffee, 81.72, 75.26, 66.50 and 43.01% when *SpliNPV* mixed with moringa + green tea, Finally it gave 71.87, 61.20, 40.62 and 24.63 % when *SpliNPV* mixed with green coffee + green tea (Table 3, Fig 3).

Table (3) Average rates of Original Activity Remaining among *S littoralis* neonate larvae treated with *SpliNPV* either alone or in combination with different antioxidants or their mixture, all exposed to different UV irradiation periods.

Irradiation exposure period (hours)	Mortality % among larvae tested with <i>SpliNPV</i> alone or mixed the indicated additives										
	<i>SpliNPV</i> alone	cacao	green tea	green coffee	moringa	cacao+ moringa	cacao + green coffee	cacao+ green tea	moringa+ green coffee	moringa + green tea	green coffee + green tea
Zero time	99.00	93.93	98.96	93.87	98.00	93.93	99.00	100.00	92.00	93.00	96.00
0.5	48.48	99.80	100.00	77.17	87.15	99.00	100.00	71.00	97.78	81.72	71.87
1	33.66	85.16	62.50	73.50	73.63	95.81	97.97	66.00	86.28	75.26	61.20
3	25.25	75.58	39.40	45.80	65.89	80.95	83.12	31.00	73.91	66.50	40.62
5	14.14	52.16	28.41	41.96	41.02	59.26	62.72	30.20	49.81	43.01	24.63




Fig (3) Average rates of Original Activity Remaining among *S littoralis* neonate larvae treated with *SpliNPV* either alone or in combination with different antioxidants or their mixture, all exposed to 300 min post UV irradiation.

Another bioassay was done with higher concentration 5% of all additives and with elongated periods of investigation which reached 10 hours. Besides the treatments that included green tea at the previous step was removed from this and further step of evaluation because it gave the lowest protection of *SpliNPV*. mortality % with *SpliNPV* alone treatment

gave 36.73, 24.74, 22.44 and 11.57 % 3, 5, 7 and 10 hours post exposure to the synthetic UV, mortality % increased to 93.87, 80.80, 70.00 and 57.00 % when *SpliNPV* mixed with cacao, 92.78, 80.61, 69.00 and 53.00 % when *SpliNPV* mixed with green coffee, 85.56, 72.44, 68.68 and 43.00 % when *SpliNPV* mixed with moringa, 97.95, 90.90, 85.56 and 59.37 % when

SpliNPV mixed with cacao + moringa, 100.00, 95.87, 90.62 and 69.47 % when *SpliNPV* mixed with cacao + green coffee, 89.79, 77.77, 71.13 and 51.04% when *SpliNPV* mixed with both moringa + green coffee. (Table 4, Fig 4) The highest LIT₅₀ found when that virus mixed with cacao + green coffee, where its activity lasts for 15.299 hours while it lasts for only 0.57 hour with virus alone treatment. Reduction % was the lowest when *SpliNPV* mixed with cacao + green coffee or

cacao + moringa where it gave only 26.53 and 39.59 %; respectively. When *SpliNPV* mixed with cacao, green coffee or moringa alone the reduction of virus activity reached 41.89, 47.00 and 57.00 %; respectively while it gave 48.96 when *SpliNPV* mixed with both moringa + green coffee, *SpliNPV* alone treatment scored the highest degree of reduction where it gave 87.41% (Table 5 & Fig 5).

Table (4) Average rates of mortality among *S littoralis* neonate larvae treated with *SpliNPV* either alone or in combination with different antioxidants or their mixture, all exposed to different further elongated UV irradiation periods.

Irradiation exposure period (hours)	Mortality % among larvae tested with <i>SpliNPV</i> alone or mixed the indicated additives at 5% total concentration						
	<i>SpliNPV</i> alone	cacao	Green coffee	moringa	cacao+ moringa	cacao + green coffee	moringa+ green coffee
Zero time	98.98 (98/99)	98.98 (98/99)	100.00 (96/96)	100.00 (96/96)	98.96 (96/97)	96.00 (96/100)	100.00 (96/96)
3	36.73 (36/98)	93.87 (92/98)	92.78 (90/97)	85.56 (83/97)	97.95 (96/98)	100.00 (98/98)	89.79 (88/98)
5	24.74 (24/97)	80.80 (80/99)	80.61 (79/98)	72.44 (71/98)	90.90 (90/99)	95.87 (93/97)	77.77 (77/99)
7	22.44 (22/98)	70.00 (70/100)	69.00 (69/100)	68.68 (68/99)	85.56 (83/97)	90.62 (87/96)	71.13 (69/97)
10	11.57 (11/95)	57.00 (57/100)	53.00 (53/100)	43.00 (43/100)	59.37 (57/96)	69.47 (66/95)	51.04 (49/96)
Control DW	0.00 (0/92)	0.00 (0/98)	0.00 (0/100)	0.00 (0/99)	0.00 (0/96)	0.00 (0/97)	0.00 (0/97)
LIT ₅₀	0.57	11.438	10.112	9.425	12.316	15.299	10.911




Fig (4) Average rates of mortality among *S littoralis* neonate larvae treated with *SpliNPV* either alone or in combination with different antioxidants or their mixtures, all exposed to 10 hours UV irradiation periods.

Table (5) Average rates of reduction among *S littoralis* neonate larvae treated with *SpliNPV* either alone or in combination with different antioxidants or their mixture, all exposed to different further elongated UV irradiation periods.

Irradiation exposure period (hours)	Mortality % among larvae tested with <i>SpliNPV</i> alone or mixed the indicated additives						
	<i>SpliNPV</i> alone	cacao	Green coffee	moringa	cacao+ moringa	cacao + green coffee	Moringa + green coffee
Zero time	98.98	98.98	100.00	100.00	98.96	96.00	100.00
3	62.52	5.11	7.22	14.44	1.01	-----	10.21
5	74.24	18.18	19.39	27.56	8.06	0.13	33.33
7	76.54	28.98	31.00	31.32	13.4	5.38	28.87
10	87.41	41.98	47.00	57.00	39.59	26.53	48.96




Fig (5) Average rates of reduction among *S littoralis* neonate larvae treated with *SpliNPV* either alone or in combination with different antioxidants or their mixture, all exposed to different prolonged UV irradiation periods.

The OAR % also gave the same trend of mortality % as showed that in (Table6 &Fig 6) where *SpliNPV* alone treatment scored 11.68 %/10 hr. post irradiation to artificial sun light while it gave 57.58,

53.00, 43.00, 59.99, 72.36 and 51.04 when *SpliNPV* mixed with cacao, green coffee, moringa or with both cacao + moringa, cacao + green coffee and moringa + green coffee. The last screening test was evaluation of additives under Egyptian sunny conditions, where *SpliNPV* alone treatment found to lose almost all its activity 48, 96 and 168 hours post irradiation where it scored 7.14, 1.03 and 1.04% of mortality only while *SpliNPV* mixed with both green

coffee and green tea treatment single out with 98.96 mortality % 168 hours post irradiation and gave almost complete protection. *SpliNPV* mixed with both cacao + moringa gave high mortality % (82.47) 96 hour post irradiation while cacao, green coffee and moringa additives gave when they mixed with *SpliNPV* 81.81, 78.12 and 74.22 mortality % 96 hours post irradiation.(table 7 & Fig 7).

Table (6) Average rates of Original Activity Remaining among *S littoralis* neonate larvae treated with *SpliNPV* either alone or in combination with different antioxidants or their mixture, all exposed to different prolonged irradiation periods.

Irradiation exposure period (hours)	Mortality % among larvae tested with <i>SpliNPV</i> alone or mixed the indicated additives						
	<i>SpliNPV</i> alone	cacao	Green coffee	moringa	cacao+ moringa	cacao + green coffee	moringa+ green coffee
Zero time	98.98	98.98	100.00	100.00	98.96	96.00	100.00
3	36.74	94.83	92.78	85.56	98.97	100.00	89.79
5	24.99	81.63	80.61	72.44	91.85	99.86	77.77
7	22.67	70.72	69.00	68.68	86.45	94.39	71.13
10	11.68	57.58	53.00	43.00	59.99	72.36	51.04




Fig (6) Average rates of Original Activity Remaining among *S littoralis* neonate larvae treated with *SpliNPV* either alone or in combination with different antioxidants or their mixture, all exposed to 10 hr. post UV irradiation.

Table (7) Average rates of mortality among *S littoralis* neonate larvae treated with *SpliNPV* either alone or in combination with different antioxidants or their mixture, all exposed to natural sunlight UV different irradiation periods.

Irradiation exposure period (hours)	Mortality % among larvae tested with <i>SpliNPV</i> alone or mixed the indicated additives at 5% total concentration					
	<i>SpliNPV</i> alone	cacao	Green coffee	moringa	cacao + green coffee	cacao+ moringa
Zero time	100.00 (99/99)	96.90 (94/97)	95.87 (93/97)	96.87 (93/96)	96.96 (96/99)	95.87 (93/97)
10	89.00 (89/100)	100.00 (98/98)	95.78 (91/95)	100.00 (96/96)	97.97 (97/99)	98.96 (96/97)
24	40.40 (40/99)	100.00 (96/96)	100.00 (96/96)	100.00 (97/97)	98.96 (96/97)	90.62 (87/96)
48	7.14 (7/98)	98.95 (95/96)	100.00 (95/95)	98.97 (97/98)	100.00 (97/97)	90.62 (87/96)
96	1.03 (1/97)	81.81 (81/99)	78.12 (75/96)	74.22 (72/97)	82.65 (81/98)	82.47 (80/97)
168	1.04 (1/96)	39.79 (39/98)	29.89 (29/97)	17.7 (17/96)	98.96 (96/97)	40.81 (40/98)
Control DW	0.00 (0/96)	0.00 (0/98)	0.00 (0/98)	0.00 (0/96)	0.00 (0/97)	0.00 (0/96)
LIT ₅₀	20.172	159.681	139.631	125.709	203.474	193.360




Fig (7) Average rates of mortality among *S littoralis* neonate larvae treated with *SpliNPV* either alone or in combination with different antioxidants or their mixtures, all exposed to 168 hours UV irradiation periods.

Reduction in virus activity was almost 100% 96 hours post irradiation where it gave 98.03 % , it scored very low rates with all additives 15.09, 17.75, 22.65, 14.31 and 13.00 % of reduction with cacao, green coffee, moringa, cacao + green coffee and cacao +

moringa additives; respectively 96 hours post irradiation. (Table 8) Virus mixed with cacao + green coffee treatment gave the highest LIT₅₀ value, where its activity lasts for 203.474 hours while it lasts for only 20.172 hours with virus alone treatment.

Original activity remaining showed that *SpliNPV* mixed with both cacao + green coffee treatment preserved 100% of its activity for five days while it gave only 1.04 % for *SpliNPV* alone treatment and gave 41.06, 31.17, 17.65 and 42.56 with cacao, green coffee, moringa and both cacao + moringa additives; respectively five days post investigation.(Table9&Fig 8)

DPPH assay

Antioxidant activity was the highest with both cacao + green coffee (85.3%) while it gave 51.03, 41.19, 33.65 and 77.56 with cacao, green coffee, moringa and both cacao + moringa additives; respectively.

Table (8) Average rates of reduction among *S littoralis* neonate larvae treated with *SpliNPV* either alone or in combination with different antioxidants or their mixture, all exposed to natural sunlight UV different irradiation periods.

Irradiation exposure period (hours)	Mortality % among larvae tested with <i>SpliNPV</i> alone or mixed the indicated additives						
	<i>SpliNPV</i> alone	cacao	Green coffee	moringa	cacao + green coffee	cacao+ moringa	
Zero time	100.00	96.90	95.87	96.87	96.96	95.87	
10	11.00	----	----	----	----	----	
24	59.60	----	----	----	----	----	
48	92.86	----	----	----	----	----	
96	98.03	15.09	17.75	22.65	14.31	13.00	

Table (9) Average rates of Original Activity Remaining among *S littoralis* neonate larvae treated with *SpliNPV* either alone or in combination with different antioxidants or their mixture, all exposed to natural sunlight UV different irradiation periods.

Irradiation exposure period (hours)	Mortality % among larvae tested with <i>SpliNPV</i> alone or mixed the indicated additives						
	<i>SpliNPV</i> alone	cacao	Green coffee	moringa	cacao + green coffee	cacao+ moringa	
Zero time	100.00	96.90	95.87	96.87	96.96	95.87	
10	89.00	100.00	99.9	100.00	100.00	100.00	
24	40.40	100.00	100.00	100.00	100.00	94.52	
48	7.14	100.00	100.00	100.00	100.00	94.52	
96	1.03	84.42	81.48	76.61	85.24	86.02	
168	1.04	41.06	31.17	17.65	100.00	42.56	




Fig (8) Average rates of Original Activity Remaining among *S littoralis* neonate larvae treated with *SpliNPV* either alone or in combination with different antioxidants or their mixture, all exposed to 168 hours post natural sunlight UV .

DISSCUTION

Baculoviruses are the major group of arthropod viruses (Herniou and Jehle, 2007). They are considered as important biological control agents (Szewczyk *et al.*, 2009), and have been used at forestry since 1900s as biological control agents, biotechnology (Contreras-Gómez *et al.*, 2014). 500 species are known of the baculovirus have been isolated from the order Lepidoptera (Possee *et al.*, 1997; Szewczyk *et al.*, 2006). As it is will known Baculoviruses, present some shortage in field stability (Stewart *et al.*, 1991). It is will proved that plant extract proved to prolong baculovirus stability against Sun light, even artificial or natural sources (Deotale *et al.*, 2007; Shapiro *et al.* 2007a, b; Shapiro *et al.*, 2008; El Salamouny *et al.*, 2009; El-Helaly *et al.*, 2009 and El-Helaly, 2013; El-Helaly *et al.*, 2013) This article provides an over view of the role of natural antioxidants and their mixtures in baculovirus protection under both natural and artificial antioxidants. It should be noted that antioxidant activity of food extracts can be determined using a (stable free radical s scavengers: galvinoxyl, diphenyl-b-picrylhydrazyl [DPPH]) (Nakatani 2003; Brown and Kelly 2007; Chen

2008; Iacopini *et al.*, 2008). These antioxidant compounds include flavonoids, phenolic acids, carotenoids, and tocopherols that can inhibit Fe3+/AA-induced oxidation, scavenges free radicals, and act as reductions (Khanduja 2003; Ozsoy *et al.*, 2009).mixing natural plants could increase oxidative stress in order to accumulative effect of antioxidants groups, both bioassay results and DPPH results proved that mixing cacao with green coffee gave highest protection to baculovirus in field and the biggest read of free radicals. This work suggest that this work should followed by studying the direct effect of different flavonoids on protection toward applicable formulation in the field.

REFERENCES

Brand-Williams W, Cuvelier ME, Berset C. 1995. Use of a free radical method to evaluate antioxidant activity. *Lebenson Wiss Technol*; 28:25-30.

Brown JE, Kelly MF. 2007. Inhibition of lipid peroxidation by anthocyanins, anthocyanidins and their phenolic degradation products. *Eur J Lipid Sci Techno* 109(1):66–71

Chen Z, 2008. Research of antioxidative capacity in essential oils of plants. *China Cond* 11:40–3.

- Contreras-Gómez AA, Sánchez-Mirón F, García-Camacho E, Molina G, Y Chisti H. (2014). Control Pestic. Sci. 51: 462-470.
- Deotale RO, Dawane PN, Biswane KD, and Borker SL. 2007. Effectiveness of UV protectants on the activity of NPV against *Helicoverpa armigera* (Hubner) on chickpea. Journal of Entomology Research. 31(1): 33-35.
- El Salamouny S, Shapiro M, Ling KS, Shepard BM. 2009. Black tea and lignin as Ultraviolet protectants for the beet armyworm nucleopolyhedrovirus. Journal of Entomological Science. 44(1): 50-58.
- El-Helaly A, El-Salamouny S, Khattab M, El-Sheikh M, Elnagar S. 2009. Preliminary evaluation of natural antioxidants as UV-protectants of *Spodoptera littoralis* nucleopolyhedrovirus, (Baculoviridae). 4th conference on recent technologies in agriculture " challenge of agriculture modernization" Cairo university, Faculty of Agriculture. 1:7-13.
- El-Helaly A, Khattab M, El-Salamouny S, El-Sheikh MA, Elnagar S. 2013. Promising additives to protect the activity of Baculovirus biocontrol agent under field – sunlight conditions in Egypt. *Journal of Life Sciences* 7: 495-500.
- El-Helaly A. 2013. Additives for a Baculovirus against Ultraviolet Effect. *Applied Science Reports*. 4 (1), 2013:187-191
- Finney DJ. 1971. Probit analysis. 3rd ed., Cambridge Univ., Cambridge U.K.
- Hemio EA, Jehle JA (2007). Baculovirus phylogeny and evolution. *Curr. Drug Targets Historical Overview and Future Outlook. Adv. Virus Res.* 68: 323-360.
- Huber J and Ludcke C. 1996. UV-inactivation of baculoviruses: the bisegmented survival curve. *Bulletin OILB/SROP*; 19(9): 253-256.
- Iacopini P, Baldi MS, Torchi P, Sebastiani L. 2008. Catechin, epicatechin, quercetin, rutin, and resveratrol in redgrapes: content, in vitro antioxidant activity and interactions. *J Food Comp Anal* 21(8):589-98.
- Khanduja KL. 2003. Stable free radical scavenging and antiperoxidative properties of resveratrol in vitro compared with some other bioflavonoids. *Ind J Biochem Biophys* 40:416-22.
- Mensor LL, Menezes FS, Leitao GG, Reis AS, dos Santos TC, Coube CS. 2001. Screening of Brazilian plant extracts for antioxidant activity by the use of DPPH free radical method. *Phyto Res*; 15: 127-130.
- Mérlillon JM, Ramawat KG (eds.) *Plant Defence: Biological Control*. Springer, Dordrecht, the Netherlands, pp: 139-160
- Mills NJ, Kean JM. 2010. Behavioral studies, molecular approaches, and modelling: 8:1043-1050.
- Moscardi F. 1999. Assessment of the application of baculoviruses for control of Lepidoptera. *Annu. methodological contributions to biological control success. Biol. Control* 52(3): 255-262.
- Muro EM, Paul JI. 1985. Laboratory evaluation of new ultraviolet absorbers for protection of Douglas-fir tussock moth (Lepidoptera: Lymantriidae) baculovirus. *Journal of Economic Entomology*. 78. 951-957.
- Nakatani N. 2003. Biologically functional constituents of spices and herbs. *J Jpn Soc Nut Food Sci* 56(6):389-95.
- Ozsoy NC, Oken EA, Kev N. 2009. Implications for degenerative disorders: antioxidative activity, total phenols, flavonoids, a scorbic acid, beta-carotene and beta-tocopherol in Aloe vera. *Oxid Med Cell Long* 2(2):99-106.
- Possee RD, Barnett AL, Hawtin RE, King LA. 1997. Engineered baculoviruses for pest Protein production using the baculovirus-insect cell expression system. *Biotechnol. Prog.* 30(1): 1-18.
- Ravensberg WJ. 2011. A road map to the successful development and commercialization of Baculoviruses-re-emerging biopesticides. *Biotechnol. Adv.* 24: 143-160
- Regnault-Roger C. 2012. Trends for commercialization of biocontrol agent (biopesticide) products. In: *Rev. Entomol.* 44: 257-289.
- Shapiro M, El Salamouny S, Shepard BM. 2008. Green tea extracts as ultraviolet protectants for the beet armyworm, *Spodoptera exigua* nucleopolyhedrovirus. *Biocontrol Science and Technology*, 18 (6): 591-603.
- Shapiro M, Shepard BM, Lopez R. 2007a. Effect of spices upon activity of gypsy moth (Lepidoptera; Lymantriidae). *Journal of Entomological Science*. 42:82-91.
- Shapiro M, Shepard BM, Lopez R. 2007b. Effect of medicinal herbs upon the biological activity of the gypsy moth nucleopolyhedrosis virus. *Biocontrol Science and Technology*. 18:605-617.
- Shorey H, Hale RL. 1965. Mass rearing of the larvae of nine noctuid species on a simple artificial medium. *Journal of Economic Entomology*. 58:522-524.
- Stewart LM, Hirst M, Ferber ML, Menyweather A, Cayley PJ, Possee RD. 1991. Construction of an improved baculovirus insecticide containing an insect-specific toxin gene. *Nature* 352: 85-88.
- Szewczyk BL, Hoyos-Carvajal M, Paluszek I, Skrzecz M, Lobo S. 2006. Baculoviruses and insect pests control in China. *Afr. J. Microbiol. Res.* 6 (2): 214-218.
- Szewczyk BL, Rabalski E, Krol W, Sihler ML. 2009. Baculovirus biopesticides—a safe alternative to chemical protection of plants. *J. Biopestic.* 2: 209-216
- Yang M, Li Y, Zhang Y, Wang J, Qu Q, Wang JY. 2012. Microbial pest control products for control of arthropods. Springer, Dordrecht, the Netherlands. pp: 171-233.

خلط مضادات الأكسدة الطبيعية يمكن أن يحسن بقاء فيروس دودة ورق القطن البوليهدروزيس النووي (Baculoviridae) تحت ظروف الحقل المصرية ألكسندرا الهاللي

قسم الحشرات الإقتصادية والمبيدات – كلية الزراعة – جامعة القاهرة

أربعة نباتات طبيعية سبق و أثبتت أنها واعدة لإطالة كفاءة فيروس دودة ورق القطن البوليهدروزيس النووي تحت الظروف المشمسة المصرية. تم اختبارها منفردة أو مخلوطة مع الفيروس في صورة أربعة مخاليط. نتائج الفحص مبنية على التقدير الحيوي، تم استخدام دلائل مختلفة لإثبات قيمة الحماية المتحصل عليها، نسبة الموت، الإختزال، الوقت اللازم للاحتفاظ بنصف الكفاءة، القدرة الفعلية ونسبة الكفاءة الحقيقية المتبقية. تم تقدير الـ (DPPH) لكل المعاملات. مخلوط الفيروس مع كل من الكاكاو + القهوة الخضراء أعطى أعلى بقاءية LIT_{50} (203.474 ساعة) بينما سجلت فقط (20.172 ساعة) مع معاملة الفيروس منفردا. الكفاءة الحقيقية المتبقية أظهرت أن مخلوط الفيروس مع كل من الكاكاو + القهوة الخضراء حفظ 100% من كفاءة الفيروس لمدة خمس أيام بينما أعطى فقط 1.04% للفيروس منفردا و أعطى 41.06، 31.17، 17.65 و 42.56 مع الكاكاو، القهوة الخضراء، المورينجا و كل من الكاكاو + المورينجا، على الترتيب خمسة أيام بعد إجراء الإختبار. هذا البحث يقترح أن خلط مضادات الأكسدة المشتقة من نباتات طبيعية توفر حماية جيدة للفيروسات العسوية المغلفة تحت الظروف المشمسة المصرية.

