

## **EFFECT OF IRRIGATION WITH DRAINAGE WATER ON PHYSIOLOGICAL PROPERTIES OF WHEAT PLANT**

**Soad S. El-Feky\* ; S. A. Gazia \*\* and Mai A. Ghazal\*\***

**\* Botany Department, Faculty of Science, Tanta University, Tanta, Egypt**

**\*\*Soil, Water and Environment Research Institute, ARC Egypt.**

### **ABSTRACT**

The impact of irrigation with wastewater supplied from drain No. 7, Kafr El-Shiekh, Egypt on wheat plant, was investigated. Growth properties, photosynthetic pigment content and the antioxidant system of wheat plant were estimated. Furthermore, heavy metals contents in the collected samples of soil, water from different sites and heavy metals accumulation in the studied plants. For this purpose, two groups of wheat plants were considered. The first group obtained from fields which irrigated using fresh water, while the other group was irrigated using water of drain No.7 from the selected five sites of drain 7. The obtained results revealed that drainage water contained significant amounts of heavy metals (Cd, Ni and Cu) warranted a pollution problem as their amounts exceed the maximum recommended concentrations according to FAO guidelines for trace metals in irrigation water. The wheat plants accumulate significant amounts of heavy metals in their shoots, straw and grains and showed a significant decrease in, fresh weight and dry weight of shoots and roots, accompanied by a marked reduction in photosynthetic pigment content. Significant increase in the activities of antioxidant enzymes and in total carbohydrate content in response to irrigation with wastewater. Cadmium was higher than critical levels at all sites in water. Copper, Cd and Pb concentration at all sites were lower than critical levels in soil. Copper, Cd and Pb concentration (ppm) in grains and straw of different sites were increased gradually from site 1 up to site 5. Lead values were the highest and Cd values were the lowest in grains and straw, but Pb concentrations at straw were higher than in grains. Copper, Cd and Pb concentrations were lower than critical levels in grains. Copper and Cd concentration were lower than critical levels, but Pb concentration was higher than critical levels in straw.

**Keywords:** Drainage Water, Irrigation, Wheat, Heavy metals and antioxidant enzymes.

### **INTRODUCTION**

Wheat (*Triticum aestivum* L.) is the most important nutritional cereal crop in Egypt. It occupies about 33% of the total winter crop area. The local production is not sufficient to meet local requirements. Egyptian production is about 7.4 million tons with an average production of 18.12 ardab per feddan (Anonymous, 2007). Food shortage and water deficit are the greatest problems discussed nowadays, and it is linked both with population growth and water allocation to different sectors, such as domestic, agronomic and industrial uses.

Egypt is mostly a rainless country that leads to limited water resources. Increasing pressure in available water resources in many areas of the world creates continued interest in use of marginal quality water such as agriculture drainage water and treated domestic wastewater. In Egypt, one example of reuse of drainage water (mixing with fresh water for irrigation) is El-Salam canal project in the North East Delta and Synaa (El-Komy, 2012). Some

researchers showed that drainage water may contain high soluble cations, anions and high EC and pH which affect on soil characters, crops growth and yield (Hendawy *et al.*, 2005; Koca *et al.*, 2007; Atlassi *et al.*, 2009 and Hamid *et al.*, 2011).

Today, the world and Egypt are facing the problem of environmental pollution. In most times drainage water contain industrial and municipal wastewater which contain heavy metals, microorganisms and parasitic (Van Breusegem *et al.*, 2001 and Koca *et al.*, 2007). Several studies have shown the effect of environmental pollutants on vegetation (Maurrary *et al.*, 1990 and Abd El Naim *et al.* 1997) and effect of heavy metals as environmental pollutants (Thomason *et al.*, 1975 and Brady, 1984).

The aim of this current study is to: investigate the effect of irrigation with wastewater supplied from different sites of drain No. 7, Kafr El-Shiekh, Egypt on wheat growth and yield production of wheat in addition to its effect on some soil characters.

## **MATERIALS AND METHODS**

### **Study area**

An experiment was conducted at Kafr El-Shiekh Governorate, Egypt using drainage and fresh water for irrigation of wheat crop in 2012 season. Drainage water was collected from different sites namely (site1 to site5) to cover all sites along away 29.39 Km from the Drain No. 7. The five sites represented drain 7 were as follow: Site (1): The beginning of the drain, Site (2): at 8.850 km, Site (3): at 12.50 km, Site (4): at 20 Km, and Site (5): Pump station of drain No. 7 which discharge the drain water into El-Burulus Lake.

Drain 7 is considered to be one of the most important drains in the Governorate of Kafr El-Shiekh with 29.390 km long. It's water used in irrigating lands of about 86.000 fed and then drained into El-Burulus Lake.

### **Sampling**

Five water samples were collected from subsurface (1.5 m) of the selected sites (site 1- site 5). The samples were taken into pre-sterilized bottles kept in iceboxes, which were further transported to the laboratory for physico-chemical analysis. For soil, samples were collected at surface level (0-30 cm in depth) from the tested five sites. The collected soil samples were air dried and sieved into coarse and fine fractions. Well mixed samples of 2 g of each fraction were taken in 250 mL glass beakers and digested with 8 mL of aqua regia on a sand bath for 2 h. After evaporation to near dryness, the samples were dissolved with 10 mL of 2% nitric acid, filtered and then diluted to 50 mL with distilled water.

### **Physico-chemical analysis**

Water temperature and pH of the collected samples were measured in the sites. Water samples were analyzed for Electrical Conductivity (EC), Dissolved Oxygen (DO), and Biological Oxygen Demand (BOD) according to standard methods (APHA, 1992).

EC of soil samples were determined in the soil paste extract by conductivity meter. Measurement of pH of the soil samples were done (soil :

water ratio 1: 2.5) were done with the help of glass electrode pH meter.

Heavy metal concentration (Cu, Cd and Pb) of each soil fraction, water samples and wheat crop (grains and straw) were assayed by Atomic Absorption Spectrophotometer using GBC. Avanta version 1.31 by flame Automization (Hanna, 1992). Quality assurance was guaranteed through double determinations and use of blanks for correction of background and other sources of error.

#### **Experimental design**

Each experiment was conducted using randomized complete blocks with three replicates. Wheat grains (*Triticum aestivum* L) were obtained from Sakha Agricultural Research Station, Kafr El-Shaikh, Egypt. Viability of wheat grains, germination percentage was estimated (95%).

#### **Growth conditions**

Wheat grains were sterilized, divided into six groups and cultivated in the selected site (near to drain No.7). One field group was irrigated with fresh water as control group. The remaining groups; were irrigated with the tested wastewater supplied from the above mentioned five sites.

Wheat grains were left to grow till the end of season, and samples (triplicates) from each group were collected at seedling (15-day old), vegetative (45-day old), flowering (70-day old) and harvesting (160-day old) stages. Each sample was subjected to the different measurements.

#### **Measurements**

##### **Growth criteria**

At vegetative and flowering stages, wheat plant were removed carefully from the cultivated soil, washed thoroughly with tap water and divided into roots and shoots. Fresh weights, dry weights and lengths of both roots and shoot were used as an indicator for studying the effect of irrigations with wastewater on wheat growth.

##### **Photosynthetic pigments**

The photosynthetic pigments; Chlorophyll a (Chl a) and Chlorophyll b (Chl b) were estimated in wheat leaves at vegetative and flowering stages as described by Arnon (1949). Briefly, 0.5 cm<sup>2</sup> of fresh leaves were homogenized immediately after harvesting in 5-10 ml aqueous acetone (85%) and kept in a dark, then centrifuged at 3000 rpm. The pigment contents were measured using a spectrophotometer at 664 and 645 nm, respectively. Final chlorophyll content (mg/g fresh. weight) was calculated according to the following equations:

$$\text{Chlorophyll a} = 10.3 E_{664} - 0.91 E_{645}$$

$$\text{Chlorophyll b} = 19.7 E_{645} - 3.8 E_{664}$$

##### **Mineral nutrient contents**

The powdered oven dry plant materials (shoots, roots and grains) were exposed to acid digestion as described by Chapman and Pratt (1961). Briefly, 0.2 g of plant sample was digested in 5 ml of H<sub>2</sub>SO<sub>4</sub> and 1 ml HClO<sub>4</sub>. The extract of digested plant materials was used for the determination of mineral nutrient contents as follow: Nitrogen (N) content by micro-Kjeldahl (Page,

1982), Total phosphorus (P) content by spectrophotometer (Snell and Snell, 1967), potassium (K) content using flame photometer (Jackson, 1967).

**Carbohydrate content:**

Carbohydrate content was determined in plant samples (shoots and grains) according to Naguib (1962). A powdered oven dry samples (0.2g) were transferred to test tubes by 10 ml 0.5 N HCl, the tubes were kept in water bath at 100°C for one hour, cooled and filtered. The filtrate was completed to 50 ml with distilled water. The reagents used were: (a) 5% phenol and (b) 95% reagent grade sulphuric acid. One ml of the extract was pipetted in a colorimetric tube. One ml of 5% phenol solution was added and mixed, 5 ml of 95% sulphuric acid was added to each tube and tubes were agitated after acid addition. After 10 minutes, the tubes were re-shaken and placed in a water bath at 25-50°C for 20 min. The absorbance was measured at 450 nm. The carbohydrate content (mg/g DW) was determined from a standard curve using glucose sugar.

**Extraction of soluble proteins**

A frozen sample (0.5 g) of wheat seedlings stage (15-day old) was homogenized in 8 ml of 50 mM cold phosphate buffer at pH 7 as described by Beauchamp and Fridovich (1971) with slight modifications. The homogenates were centrifuged at 4000 rpm for 20 min, the collected supernatant was used as a crude extract for the following antioxidant enzymes assay:

**1- Catalase**

Catalase was assayed by measuring the initial rate of disappearance of H<sub>2</sub>O<sub>2</sub> (Kato and Shimizu, 1987). A sample of 3 ml of reaction mixture contained 0.1 M sodium phosphate buffer (pH 7), 2 mM H<sub>2</sub>O<sub>2</sub> and 0.1 ml enzyme extract. The decrease in H<sub>2</sub>O<sub>2</sub> was followed as a decline in the absorbance at 240 nm. The activity was expressed in units of  $\mu$ M of the substrate converted per minute per gram fresh weight.

**2- Peroxidase**

Peroxidase activity was measured according to Kato and Shimizu, (1987). The assay medium contained 0.1 M sodium phosphate buffer (pH 5.8), 7.2 mM guaiacol, 11.8 mM H<sub>2</sub>O<sub>2</sub> and 0.1 ml enzyme extract and the final assay volume was 3 ml. The reaction was initiated by the addition of H<sub>2</sub>O<sub>2</sub> and the change in absorbance was at 470 nm. Enzyme activity was expressed in units of  $\mu$ M of the substrate converted per min per gram fresh weight.

**3- Superoxide dismutase**

Superoxide dismutase was assayed on the basis of its ability to inhibit the photochemical reduction of nitro blue tetrazolium (Beauchamp and Fridovich, 1971). Three ml of the reaction mixture contained 50 mM sodium phosphate buffer (pH 7.8), 13 mM methionine, 75 mM nitro blue tetrazolium, 2  $\mu$ M riboflavin, 100 mM EDTA and 0-200  $\mu$ l enzyme extract. Absorbance by the reaction mixture was at 560 nm.

**Statistical analysis**

All the obtained data was statistically analyzed using the methods described by Cochran and Cox (1960).

## RESULTS AND DISCUSSION

According to the FAO Land and Plant Nutrition Management Service, over 6% of the world's land is affected by either salinity or sodicity. Moreover, the low water quality and the poor drainage systems are the greatest causes of these stresses, and this problem is more acute with higher evaporation, especially in arid and semi-arid zones. Use of industrial or municipal waste water for irrigation purpose is a common practice (Singh *et al.*, 2004). The effluent not only affects the plant growth but also deteriorate the soil properties when used for irrigation where saline soils are wide spread that induced the decreasing of land productivity in many countries over the world (Atlassi *et al.*, 2009).

### Physico-chemical analysis

Resulte in Table (1) showed the variation of EC, Na<sup>+</sup>, Ca<sup>++</sup>, Mg<sup>++</sup>, K<sup>+</sup>, HCO<sub>3</sub><sup>-</sup> and Cl<sup>-</sup> values at different five sites along drain No.7. The values of Na<sup>+</sup>, Ca<sup>++</sup>, Mg<sup>++</sup>, HCO<sub>3</sub> and Cl were heigher than ISI ( standard for irrigation). The lowest values of EC, Na<sup>+</sup>, Ca<sup>++</sup>, Mg<sup>++</sup>, K<sup>+</sup>, HCO<sub>3</sub><sup>-</sup> and Cl<sup>-</sup> were recorded at site 1 (0.82, 5.6, 1.3, 1.8, 0.1, 3.5 and 3.9 ) respectively, while the heighest values were recorded at site 2 (5.58, 37.6, 8.9, 12.3, 0.6, 0.6 and 26.6 ) respectively.

This may be de to soluble salts discharged to the drain from the soils surrounded by the drain throught the drainage or soluble salts from municiple wastewater, industerial water discharged to the drain No.7.This resulte are agreed with those obtained by El-Maghraby, (2008).

**Table 1. Chemical characteristics of irrigation water from different sites of drain No. 7 :**

Water analysis	site1	site2	site3	site4	site5	IS standards for irrigation (ISI)
EC dSm <sup>-1</sup>	0.82	5.58	1.47	1.71	4.16	0.42
Na <sup>+</sup> meq/l	5.6	37.6	10	11.6	28.3	1.3
Ca <sup>++</sup> meq/l	1.3	8.9	2.4	2.4	6.7	1.7
Mg <sup>++</sup> meq/l	1.8	12.3	3.2	3.2	9.2	0.97
K <sup>+</sup> meq/l	0.1	0.6	0.1	0.1	0.4	0.23
CO <sub>3</sub> <sup>-</sup> meq/l	0	0	0	0	0	0
HCO <sub>3</sub> <sup>-</sup> meq/l	3.5	0.6	7	6.5	6	3.09
Cl meq/l	3.9	26.6	0.7	8.1	19.9	0.75

### Chemical characteristics of soil from drain No.7 :

Table (2) showed that the variation of EC, Na<sup>+</sup>, Ca<sup>++</sup>, Mg<sup>++</sup>, K<sup>+</sup>, CO<sub>3</sub>, HCO<sub>3</sub><sup>-</sup> and Cl<sup>-</sup> values at different five sites of soil from drain No.7. in soil of different locations at drain no. 7 The lowest values of EC, Na<sup>+</sup>, Ca<sup>++</sup>, Mg<sup>++</sup>, K<sup>+</sup>, CO<sub>3</sub>, HCO<sub>3</sub><sup>-</sup> and Cl<sup>-</sup> were recorded at site 3 (1, 6.8, 1.6, 2.2, 0.1, 0, 7 and 4.8 ) respectively, while the heighest values were recorded at site 5 (4.5, 30.6, 7.2, 9.9, 0.5, 0, 9 and 21.4), respectively, followed by site 4 and the lowest site was site 3. EC at site 5 were the heighest. These resultes are

relatively in complete accordance with the results which recorded by Ladwani *et al.*, (2012) and Mir *et al.*, (2012) who they reported that the untreated or partially treated waste water from industries are continuously used in irrigating the agricultural fields in developing countries and continuous use of this waste water for irrigation effects of soil quality.

**Table 2. Chemical characteristics of soil samples from different sites of drain No. 7 :**

Water analysis	site1	site2	site3	site4	site5	IS standards for irrigation (ISI)
EC dSm <sup>-1</sup>	2.25	2.11	1	3.9	4.5	5.0
Na <sup>+</sup> meq/l	15.3	14.3	6.8	26.5	30.6	14.8
Ca <sup>++</sup> meq/l	3.6	3.4	1.6	6.2	7.2	21.2
Mg <sup>++</sup> meq/l	5	4.6	2.2	8.6	9.9	10.4
K <sup>+</sup> meq/l	0.2	0.2	0.1	0.4	0.5	0.3
CO <sub>3</sub> <sup>--</sup> meq/l	0	0	0	0	0	0.0
HCO <sub>3</sub> <sup>-</sup> meq/l	7	8	7	7	9	2.8
Cl <sup>-</sup> meq/l	10.7	10	4.8	18.6	21.4	10.8

Heavy metals (Cu, Cd and Pb) concentrations (mg/L) were estimated in the collected wastewater and soil samples at different sites (site.1-site.5) as represented in Table 3. The obtained data revealed that the highest values of the three heavy metals were recorded in site.5 and the lowest values recorded in site.1. As observed, Pb concentrations ranged between 4.6; (site1) to 6.3; (site4) and 0.075; (site 1) to 1.5; site.4 ppm in soil and wastewater samples, respectively. On other hand Cd concentrations recorded the lowest values in collected soil and wastewater samples. The highest value (.119 ppm) of Cu concentration was estimated in water samples collected from site.5 and the lowest value (0.036) was detected in site 1.

The increases of the studied three heavy metals in samples of site 1 ) may be due to that site 1 is near from the first drainage source. Also the three heavy metals were increased gradually due increases of heavy metals leached from the soils during reuse of the drainage water. These results agreement with El-Guindy and Amer, (1989) who reported that trace elements occur in almost all drainage water but at low concentrations, Mohamed *et al.*, (1998) detected that in spite of the low concentration of heavy metals in irrigation water, high accumulation of Pb and Cu were observed in the soil and crop yields, Alnenaie (2003) stated that, the contamination of water bodies in the Egyptian Nile Delta by metals is caused by the discharge of massive amounts of domestic sewage as well as agricultural and industrial effluents, Franca *et al.*, (2005 ) observed that heavy metals occur in the environment both as a result of natural processes and as pollutants through various pollution sources from human activities. Maceda-Veiga *et al.*, (2012) detected that domestic wastewaters can contain fairly high concentrations of metals, such as Al, Ni, Cu, Fe, Pb and Zn, which are derived from a wide variety of household products, such as cleaning materials, toothpaste and cosmetics. Ghannam *et al.*, (2014) concluded that Cd was not detectable (below the detection limits with ICP -OES <0.001 and <0.005) in water

samples ( El-Bahr El-Pharaony drain). On the other hand, Cu and Pb exceeded the permissible limits of Egyptian law.

The increases of the studied three heavy metals (Cu, Cd, Pb) from site 1 up to site 5 in soil of different locations at drain 7 may be due to using water of drain 7 for irrigation which precipitate heavy metals in the soil. Results are relatively in complete accordance with the results recorded by Nriagn and Pacyna, (1988) who reported that about 2 million tons of copper are released each year as a result of human activity, causing locally raised levels in terrestrial and aquatic environments, Franca *et al.*, (2005) observed that heavy metals occur in the environment both as a result of natural processes and as pollutants through various pollution sources from human activities. And Ladwani *et al.*, (2012) and Mir *et al.*, (2012) so they reported that the untreated or partially treated waste water from industries are continuously used in irrigating the agricultural fields in developing countries and continuous use of this waste water for irrigation effects of soil quality.

**Table (3) Heavy metals content (ppm) of different waste water and soil sample collected from different sites of drain No.7.**

Collected samples from different sites	Soil			Water		
	Cu	Cd	Pb	Cu	Cd	Pb
Site.1	3.34±0.314	0.108 ±0.011	4.3±0.2	0.036±0.006	0.027±0.097	1.102±0.097
Site.2	3.65±0.216	0.082±0.011**	4.63±0.157*	0.039±0.001	0.082±0.011**	1.05±0.010
Site.3	5.49 ±0.012**	0.089±0.009**	5.23±0.208**	0.12±0.01**	0.089±0.009**	1.306 ±0.015**
Site.4	8.02±0.020**	0.132±0.006**	5.6±1.087**	0.112±0.001**	0.132±0.006**	1.484±0.014**
Site.5	8.83 ±0.079**	0.139±0.00**	6.33±0.152**	0.119±0.001**	0.139±0.00**	1.532±0.010**
ISI	0.05	0.0991	3.97	0.015	0.005	0.075
L.S.D <sub>0.05</sub>	0.317	0.015	0.293	0.010	0.015	0.081
L.S.D <sub>0.01</sub>	0.451	0.021	0.417	0.014	0.021	0.115

**(ISI) Is Standards for Irrigation**

**Critical levels in soil according to Kabata –Pendias and Pendias (1992) for Cu, Cd and Pb were 60 : 125, 3 : 8 and 100 : 400 ppm respectively.**

**Critical levels in water according to FAO (1985) for Cu, Cd and Pb were 0.2, 0.01 and 5 ppm, respectively.**

**\* Results are significantly different from control at (p<0.05)**

**\*\* Results are significantly different from control at (p<0.01).**

### **Effect of irrigations with wastewater on growth of wheat**

Growth criteria of wheat plant irrigated with water from the different sites of drain No.7, at vegetative and flowering stages were measured. The obtained results presented in Fig (1) indicated that the maximum growth of wheat plants was recorded in group irrigated with fresh water (control) rather than the wastewater (site 1- site5). Shoot fresh weight at vegetative and flowering stages of plant samples were significantly decreased in comparing with control samples (1.2-2.8 g vs 3.0 g and 0.06 - 0.6 g vs 0.9, respectively). Noticely, site 5 showed the upper most deleterious effect on growth criteria obtained from wheat crop. These results are agreed with those obtained by

**Soad S. El-Feky et al.**

Atlassi *et al.*, ( 2009). Heandway *et al.*, (2005) and Koca *et al.*, 2007 who stated that , waste water and polluted soil decreases in growth of crops, may be due to toxicity of high level of dissolved solids. The increase of shoot and root fresh weight of samples irrigated with water from site 3 may be due to discharged of drainage water less in EC and soluble cations and anions.


**Fig. 1 . Growth criteria of wheat plant groups at vegetative and flowering stages.**



**Effect of irrigations with wastewater on chlorophyll content of wheat**

The chlorophyll (a and b) content in wheat shoots of all wheat groups were presented in Fig 2. The data indicated that chlorophyll content in wheat plants irrigated with fresh water recorded higher values than those irrigated by wastewater at both vegetative and flowering stages. Chl a and Chl b content of wheat plant irrigated with wastewater from site1 at vegetative stage was decreased with about 7.5% and 1.82%, respectively with comparing to control group . While, at flowering stage the decrease of Chl a and Chl b content was about 2.65% and 3.37% respectively, compared to control. However, samples from site 5 showed a highly significant decrease in Chl a and Chl b content (38% and 36.36%, respectively) at vegetative stage while it reached (39.07%) and (38.65%) reduction at flowering stage. Enhancement of chlorophyll content in both wheat samples irrigated with water from site 1 and control samples may be due to higher nutrient uptake from the water. Nagda *et al.*, (2006) in their experiment on seed germination found similar result. Srivastava and Sanhai (1987) also found similar results using distillery wastewater. Higher concentration of wastewater are inhibitory to synthesis of chlorophyll molecules particularly chlorophyll a (Khan *et al.*, 2011). The variation in chlorophyll content was found to be statistically significant with treatment at various sites of drain No. 7.

The change in chlorophyll content indicates that the chlorophyll synthesizing capacity have diminished affecting of the photosynthetic process. Krupa *et al.*, (1993) reported values in the same trends as we were investigated, they mentioned that cadmium content in wastewater reduced plant growth, biomass and leaf pigments, and cadmium can harness photosynthetic activity as well as the chlorophyll content.



**Fig. 2. Chlorophyll a and b content on wheat crop groups at vegetative and flowering stages.**

**Effect of irrigations with wastewater on total carbohydrates % of wheat**

Total carbohydrate (mg\gm DW) in shoot at flowering stage of wheat plant groups (G1-G5) were increased than those of control samples as shown

in Table 4. At flowering stage of wheat plant G5, were increased by 37.9%, 8.2% and 27.8 % in shoot, grains and straw, respectively compared with control samples. The lowest values of carbohydrate content was recorded samples irrigated with water from site 1, however, total carbohydrate in grains and straw at harvest stage decreased significantly with about (1.7%) and (1.4%) respectively.

This may be due to stress effect which reflected on wheat leaf area. Carbohydrate content in straw and grain at harvest stage (Table 4) were decreased in samples obtained from site 1, this may be due to leaf area was less than plants which irrigated by drainage water comparing with control which irrigated by fresh water so effect on photosynthesis and carbohydrate accumulation. These results are harmony with those obtained by Rong *et al.*, (2007) who showed that stress induces the decline in protein contents in plants but increase in soluble sugar content. Several studies suggested that the accumulation of soluble carbohydrates was significantly related to salt tolerance (Teimouria *et al.*, 2009; Yin *et al.*, 2010 and Zhang *et al.*, 2012). Hirich *et al.*, 2014, stated that soluble sugars increased with increasing irrigation water salinity.

**Table 4 . Total carbohydrate (mg/gm DW) in shoot, grains and straw, of wheat plants irrigated by water from different locations of drain No. 7.**

Wheat crop group	shoot	grains	straw
control	62.496± 2.739	74.837±12.32	65.879±3.733
Site 1	66.117±.818*	73.577±2.769	64.947±2.868
Site2	81.370±1.516**	80.0±0**	68.290±2.211
Site3	75.122 ±3.530**	55.943±0.863**	62.097±2.539
Site 4	85.694±.795**	85.057±2.280**	78.643±1.794**
Site 5	86.182 ±.743**	88.470±1.371 **	84.190±2.786**
L.S.D 0.05	3.566	2.995	4.843
L.S.D 0.01	5.000	4.199	6.790

\*Results are significantly different from control at (p <0.05)

\*\*Results are significantly different from control at (p <0.01).

Each result is triplicate reading

#### Heavy metals concentration in straw and grains of wheat

Heavy metals (Cu, Cd, Pb) content in straw and grains of wheat plant irrigated with water collected from different station in drain No.7 are shown in (Table 5).

Concentrations of three heavy metals (Cu, Cd and Pb) ppm in grains and straw at harvest stage of samples irrigated with water from the different sites of drain 7 increased significantly than those of the control samples. Meanwhile samples of the site 1 showed the lowest concentration. Three heavy metals (Cu, Cd, and pb ) in grains of site1 was increased with about (42.8%), (103.3%) and (325.6%), respectively. However, site 5 showed a highly significant increase were about (126.1%), (155.9%) and (545.83%), respectively, compared to control treatment.

However, evolution for three heavy metal (Cu, Cd and pb ) in straw of site 1was increased significantly with about (20.9%), (5.1%) and (223%)

respectively. And of site 5 showed a highly significant increase was about (112.4%), (106.3%) and (263.6%), respectively, compared to control treatment. Pb concentration in grains and straw were higher than Cu and Cd concentration. While concentrations of (Cu, Cd and Pb) in straw are higher than in grains. Pb concentration in straw were higher than critical levels in according to FAO (1985) .

The increases of the studied three heavy metals (Cu, Cd, Pb) in straw and grains of wheat plant which irrigated with water of different sites from drain No.7 (Table 5) may be due to the presence of these heavy metals in the water and the soil, similar results were obtained by Pietz *et al.*, (1978) and Amin, (1997) who reported that Pb concentration in all studied crop irrigated with fresh and drainage water was higher than the permissible levels for human consumption.

**Table 5 . Heavy metals contents (mg/ Kg) of grains and straw samples at harvest stages from wheat plant.**

Water used for irrigations collected from different sites	Collected samples of wheat irrigated by wastewater					
	Grains			Straw		
	Cu	Cd	Pb	Cu	Cd	Pb
CG	0.84±0.05	0.059±0.01	0.336±0.208	6.78±0.243	0.253±0.0585	5.5±0.2
St.1	1.2±0.01**	0.12±0.01**	1.43±0.09**	8.2±0.1	0.24±0.01	17.8±0.328
St.2	1.28±0.026 **	0.34±.01**	3.21±0.05**	9.5±0.141**	0.525±0.005**	18.4±0.1**
St.3	1.5±0.1**	0.12±0.01**	1.67±0.49**	10.46±0.057**	0.336±0.025**	18.5±0.0**
St.4	1.73±0.01**	0.176±0.015**	2.64±0.17**	13.2±0.1**	0.736±0.015**	19.7±0.132**
St. 5	1.9±0.055**	0.151±0.005**	2.17±0.06**	14.4±0.1**	0.522±0.006**	20±0.0577**
L.S.D <sub>0.05</sub>	0.118	0.018	0.423	0.243	0.048	0.307
L.S.D <sub>0.01</sub>	0.165	0.026	0.593	0.341	0.068	0.43

**Critical levels in grains and straw according to FAO (1985) for Cu, Cd and Pb were 4:15, 0.2:0.8 and 4:15 ppm respectively.**

**\* Results are significantly different from control at (p<0.05)**

**\*\* Results are significantly different from control at (p<0.01)**

### **Effect of irrigations with wastewater in antioxidant enzymes produced in wheat seedlings**

The activities of antioxidant enzymes Catalase (CAT), Peroxidase (APX) and Superoxide Dismutase (SOD) in leaves of wheat plant irrigated by wastewater (G1 to G5) are shown in Fig. (3). Antioxidant enzymes activities ( $\mu$  l / g f.w) of wheat samples groups irrigated with wastewater were increased significantly than those of CG. Meanwhile, catalase, SOD and APX in G1 samples were increased than CG by 16%, 19.3% and 11.7%, respectively. Samples of G5 has the highest increase with 88.2 %, 69.1% and 64.7 % for catalase, peroxidase and SOD, respectively.

Drainage water contains salts and heavy metals in varied degrees led to oxidative stress in plant irrigated with water and cause several physiological changes due to change in metabolic pathway (Lee *et al.*, 2001;

Panda and Upadhyay, 2003). Production of several Active Oxygen Species (AOS) increases in the presence of NaCl and has been stated to damage almost every macromolecule (Khan and Panda, 2002). However, in many plant cells exposed to oxidative stress the lead to induction of both enzymatic (Superoxide Dismutase (SOD), Catalase (CAT).

Changes of antioxidant enzymes activity under environmental stresses have been suggested as indicators for biotic or abiotic stresses (Lee, 1997; Pakniyat and Abedi 2010 and Tian *et al.*, 2012).

**Fig. 3. Effect of irrigation with water from different sites on antioxidant enzymes activities (catalase, super oxide dismutase, and peroxidase).**

- This it can be concluded that water from site 1 have the lower deliterius effect so we can use it in irrigation, however water from site 5 have the most deliterius effect so we must not use it in irrigation.

## **REFERENCES**

- Abd El-Naim, M., El-Mowelhi, N. and El-Houseini, M. (1997). Risk assessment of sewage sludge application on agricultural land in Egypt. International Symposium on Salt Affected Soil. Cairo 22-26 Sept. 1997.pp-:523-574.
- Alnenaï, A.A. (2003). Contamination of irrigation and drainage canals and ponds in the Nile Delta by heavy metals and its association with human health risks. Egypt. J. Zoology, 41: 47-60.
- American Public Health Association (APHA) (1992). Stander methods for examination of water, and wastewater methods for examination of water, and wastewater<sup>18<sup>th</sup></sup>.edn. Washington.D.C.

- Amin, M. Sh. (1997). Effect of irrigation with wastewater and N,P & K fertilization on production and chemical composition of wheat and maize grain under sandy soil condition. M. Sc. Thesis Fac. Agric. Cairo Univ. (1997).
- Anonymous (2007). Wheat area and production. Wheat Research Center, Field Crops Research Institute ARC, Ministry of Agriculture & land Reclamation.
- Arnon, D. I. (1949). Copper enzymes in isolated chloroplasts. Polyphenol oxidase in *Beta vulgaris*. Plant physiology. 24:1-15.
- Atlassi P.V., Nabipour M. and Meskarbashee, M. (2009): Effect of salt stress on chlorophyll content, Fluorescence, Na<sup>+</sup> and K<sup>+</sup> ions content in Rape plants (*Brassica napus* L.). Asian Journal of Agriculture, 3(2): 28-37.
- Beauchamp, c. and Fridovich, I. (1971). Super oxide dismutase: improved assays and an assays applicable to acrylamide gels .Analytical Biochemistry .44:276-287.
- Brady, N. (1984). The Nature and Property of Soil (New York: Macmillan).
- Chapman, H.D. and pratt, P.F. (1961). "Methods of Analysis for Soils, Plant and Water". University of California, agriculture publication. Barkely, California, PP.17.
- Cochran, W.G. and Cox, G.M. (1960). Experimental Designs, 2<sup>nd</sup> ed. John Welly, New York, PP.293-316.
- El-Guindy, S. and Amer, M.H. (1989). Quality of drainage water in the Nile Delta. Toxic substances in agricultural water supply and conference irrigation and drainage. Ottawa, Canda, June. 8-9, pp 351-366.
- El-Komy, M. N. A. E. (2012). Assessing the quality of irrigation water resources in north Kafr-El-Sheikh governorate .M.Sci.Thises. Fac. Agric.Mansoura Univ.
- El-Maghraby, F. M.I. (2008). Studies on Non- Symbiotic nitrogen fixing bacteria in Agricultural Drainage water Drain (7) in Kafr El-Sheikh Governorate. MSc 2008. Faculty of Science. Tanta University.
- FAO. (1985). Water quality for agriculture. R.S. Ayers and D.W. Westcot. Irrigation and Drainage Paper 29 Rev. 1. FAO, Rome. 174 p.
- Franca, S.; Vinagra, C.; Cacada, I. and Cabral, H.N. (2005). Heavy metal concentrations in sediment, benthic invertebrates and fish in these salt marsh areas subjected to different pollution loads in the Tagus Estuary (Portugal). Mar. Pollut. Bull., 50: 997-1003.
- Ghannam, Hala, E. ; Talab, S. A. ; Jahin, H. S. and Gaber, S. E. (2014). Seasonal variations in physicochemical parameters and heavy metals in water of El-Bahr El-Pharaony Drain, El-Menoufia Governorate, Egypt. Research Journal of Environmental and Earth Sciences 6(3): 174-181.
- Hamid, M. Mostafa, H. and Mohammad, A. (2011). Physiological and biochemical responses of wheat cultivars under salinity stress. ARPN Journal of Agricultural and Biological Science. VOL. 6, NO. 5.

- Hanna, R. G. (1992). Sequential extraction metals in cooks River sediments. Reprint from: Miskiewicz., A.G.(ed.) (1992). Proceedings of a Bioaccumulation of contaminants in Aquatic Environments. Water Board and Australian Marine Sciences Association. Inc., Sydney.
- Hendawy S., Elhua Y., Yakout G.M., Awad A.M., Hafiz S.E. and Schmidhalter U. (2005). Evaluating salt tolerance of wheat genotypes using multiple parameters. *Europ. J. Agronomy*, 22: 243-253.
- Hirich ,A. ; El Omari, H. ; Jacobsen, S. ; Lamaddalena, N. ; Hamdy, A. ; Ragab, R.; Jelloul, A., and Choukr-Allah, R. (2014). Chickpea (*Cicer arietinum* L.) physiological, chemical and growth responses to irrigation with saline water. *Australian Journal of Crop Science*. AJCS 8(5):646-654.
- Jackson, M.L. (1967). "Soil Chemical Analysis". Prentice Hall of India, private limited, New DeLhi .
- Kabata Pendias, A. and Pendias, H. (1992). "Trace Elements in Soils and Plants". (2nd ed) CRC press, Boca Raton, Flo. USA.
- Kato, M. and Shimizu, S. (1987). Chlorophyll metabolism in nigher plants .VII. Chlorophyll degradation in senescing tabacco leaves; phenolic-dependent peroxidase degradation .*Condian Journal of Botany*.65: 729-735.
- Khan, M.H. and Panda, S.K. (2002). Induction of oxidative stress in roots of oryza sativa l. in response to salt stress. *Biol. Plantarum*, 45: 625-627.
- Khan, M.G.; Danlel, G.; Konjit, M.; Thomas, A.; Eyasu, S.S. and Awoke, G. (2011). Impact of textile waste water on seed germination and some physiological parameters in pea (*Pisum sativum* L.), Lentil (*Lens esculentum* L.) and gram (*Cicer arietinum* L.). *Asian Journal of Plant Science*, 10, 269-273.
- Krupa ,Z.; Oquist, G. and Huner, N. P. A. (1993). The effect of cadmium on photosynthesis of *Phaseolus vulgaris* L. a fluorescence analysis. *Physio.plantarum* 88, 626- 630.
- Koca M.; Bor, M.; Ozdemir, F. and Turkan, I.(2007). The effect of salt stress on lipid peroxidation, antioxidative enzymes and proline content of sesame cultivars. *Environ Exp. Bot.* 60: 344-351.
- Ladwani, K.D.; Ladwani, K. D.; Manik V. S. and Ramteke, D. S. (2012). Impact of Industrial Effluent Discharge on Physico-Chemical Characteristics of Agricultural Soil, *International Res.J. Of Env.Sci*, 1(3), 32- 36.
- Lee, M.Y. (1997). The changes of peroxidase activity and isoperoxidase patterns from pine needles under the salinary stress *The Korean Journal of Ecology* 20(5) :315-321.
- Lee, H.D.; Kim, Y.S. and Lee, C.B. (2001). The inductive responses of the antioxidant enzymes by salt stress in the rice (*Oryza sativa* L.). *J. Plant Physiol.*, 158:737-745.
- Maceda-Veiga, A.; Monroy, M. and de Sostoa, A. (2012). Metal bioaccumulation in the Mediterranean barbel (*Barbus meridionalis*) in a Mediterranean River receiving effluents from urban and industrial wastewater treatment plants. *Ecotox. Environ. Safe.* 76: 93-101.

- Murray, A. N.; Gupta, R.K.; Usha, G. M. and Dhingra, N. S. (1990). Effect of coal smoke pollutants of iron foundries on fishes religiose line. Indian J. Environ. Health, 32, 163-166
- Mir, T. A. ; Manderia, S. and Manderia, K..(2012). Influence of dye industrial effluent on physico chemical characteristics properties of soil at Bhairavgarh, Ujjain, MP, India, International Res.J. of Env.Sci, 1(1), 50-53.
- Mohamed, A.; Rashad, A.; Soheir, F., and Ahmed, S.(1998). Risk assessment of soil and water pollution on agricultural crops in Egypt. Agricultural Research Center, 9 El-Gamaa str. Giza, Egypt.
- Nagda, G.K.; Diwan, A.M. and Ghole, V.S. (2006). Seed germination bioassays to assess toxicity of molasses fermentation based bulk drug industry effluent. Electronic Journal of Environmental, Agricultural and Food Chemistry 5(6), 1598-1603.
- Naguib, M. I. (1962). Arapid colorimetric procedure for the estimation of free and conjugated sugars in plant extracts .Zucker. 15:351.
- Nriagn, J.O. and Pacyna, J. M. (1988). Quantitative assessment of worldwide contamination of air, water and soil by trace metals. Nature 333, pp 134-139.
- Page, A. L.; Miller, H. and Keeney, D. R. (1982)." Methods of Soil Analysis". Part 2, ASA, SSSA, Madison, Wisconsin USA.
- Panda, S.K. and Upadhyay, R.K. (2003). Salt stress injury induces oxidative alterations and antioxidative defiance in the roots of Lemna minor. Biol. Plantarum, 48: 249-253.
- Pakniyat, H. and Abedi, T. (2010). Antioxidant enzyme changes in response to drought stress in ten cultivars of oilseed rape (Brassica napus L.) Czech J. Genet. Plant Breed. 46 (1): 27-34.
- Pietz, R.; Vetter, I. ; Masarik, D. and Mcfree, W. W. (1978). Zinc and Cadmium contents of agricultural soils and corn in north western Indiana. Environ. Qual 7, 381-385.
- Rong, G. T. ; Guo P. Z. and Yan, H. Z. (2007). Physiological changes in barley plants under combined toxicity of aluminum, copper and cadmium. Colloids and Surfaces. B. Biointerfaces, 57: 182-188.
- Singh, K.P., Mohem, D., Sinha, S., and Dalwani, R. (2004). Impact assessment of treated/untreated waste water toxicants discharged by sewage treatment plants on health, agriculture and environmental quality in the waste water disposal area. *Chemosphere*, 55: 227-255.
- Snell, F.D. and Snell, C.T. (1967). Colorimetric methods of analysis .Condian Journal of Botany.63: 729-735.
- Srivastava, N. and Sanhai, R. (1987). Effects of distillery waste in the performance of *Cicer arietinum* L. Environmental pollution 43, 91-102.
- Teimouria, A.; Jafarib, M. and Azarnivand, H. (2009). Effect of proline, soluble carbohydrates and water potential on resistance to salinity of three Salsola species (*S.rigida*, *S. dendroides*, *S.richter*). Desert 14:15-20.

- Thomason, B. M.; Biddle, J. W. and Cherry, W. B. (1975). Detection of salmonellae in the environment. Journal of Applied Microbiology; 30(5): 764-767.
- Tian, Z.; Wang, F.; Zhang, W.; Liu, C. and Zhao, X. (2012). Antioxidant mechanism and lipid peroxidation patterns in leaves and petals of marigold in response to drought stress. Hortic. Environ. Biotechnol., 53(3):183-192.
- Van Breusegem F.; Vranova, E.; Dat, J. F. and Inze, D. (2001). The role of active oxygen species in plant signal transduction. Plant Sci. 161: 405-414.
- Yin, Y.G. ; Kobayashi, Y.; Sanuki, A.;Kondo, S. ; Fukuda, N.; Ezura, H.; Sugaya, S. and Matsukura, C. (2010). Salinity induces carbohydrate accumulation and sugar-regulated starch biosynthetic genes in tomato (*Solanum lycopersicum* L. cv. 'Micro-Tom') fruits in an ABA- and osmotic stress-independent manner. J Exp Bot 61:563-574.
- Zhang-hua, Z.; Qiang, L.; Hai-xing1, S.; Xiang-min, R. and Ismail, A.M. (2012). Responses of different rice (*Oryza sativa* L.) genotypes to salt stress and relation to carbohydrate metabolism and chlorophyll content. Afr J Agri Res 7:19-27.

### تأثير الري بمياه الصرف الصحي على الخواص الفسيولوجية لنبات القمح

سعاد سليمان الفقي\*، السيد عامر جازية\*\* و مى عبد اللطيف غزال\*\*  
\* قسم النبات - كلية العلوم - جامعة طنطا - مصر  
\*\* معهد بحوث الأراضى والمياه والبيئة - مركز البحوث الزراعية - مصر

تم دراسة تأثير ري نباتات القمح بمياه مصرف نمره ٧ بمحافظة كفر الشيخ وقدرت مقاييس النمو والمحتوى الصيغى ونظام مضادات الأكسدة لنبات القمح بالإضافة إلى ذلك قدر محتوى عينات التربة والنبات من العناصر الثقيلة .

ولتحقيق هذا الغرض تم أخذ مجموعتان من عينات القمح : المجموعة الأولى أخذت من حقول تروى بماء عذب بينما المجموعة الأخرى تروى بمياه مصرف نمره ٧ من الخمس مواقع المختاره وقد أظهرت النتائج ان مياه الصرف تحتوى على كميات من العناصر الثقيلة (كادميوم ورسااص ونحاس) مما يتسبب فى مشكلة التلوث إذا ما تعدت الكميات الموجودة فى مياه الري الحدود الموصى بها حسب إرشادات منظمة الأغذية والزراعة.

وقد تجمعت كميات ملحوظة من العناصر الثقيلة فى كل من الجزء الخضرى وقش وحبوب القمح مما أدى إلى نقص ملحوظ فى الوزن الرطب والوزن الجاف لكل من الجذور والجزء الخضرى مصحوبة بنقص فى المحتوى الصيغى وكذلك زيادة ملحوظة فى نشاط الإنزيمات المضادة للأكسدة وكذا المحتوى الكلى للكربوهيدرات إستجابة للرى بمياه الصرف.

سجل عنصر الكادميوم فى عينات المياه نتائج أعلى من الحد الحرج فى كل المواقع بينما كان مستوى النحاس والكادميوم والرسااص فى التربة أقل من الحد الحرج وزاد تركيز النحاس والكادميوم والرسااص فى حبوب وقش نبات القمح تدريجياً من الموقع (١) الى الموقع (٥).

كان أعلى العناصر تركيزاً هو الرسااص وأقلها الكادميوم فى كل من الحبوب والقش وكان الرسااص فى القش أعلى من الحبوب.

بالنسبة للحبوب كان تركيز النحاس والكادميوم والرسااص أقل من الحد الحرج وفى القش كان تركيز النحاس والكادميوم أقل من الحد الحرج بينما تركيز الرسااص كان أعلى من الحد الحرج.