

IDENTIFICATION AND CHARACTERIZATION OF ACTINOMYCETES FOR BIOLOGICAL CONTROL OF BACTERIAL SCAB OF *STREPTOMYCES SCABIES* ISOLATED FROM POTATO

Hind A. Hamedo⁽¹⁾ and Abeer H. Makhoul⁽²⁾

⁽¹⁾ Faculty of Education, El Arish Suez Canal University

⁽²⁾ Faculty of Agriculture, Minufiya University

(Received: Sep. 30, 2012)

ABSTRACT: A novel strain of *Streptomyces* in Egypt (known as *Streptomyces scabies*) was isolated from a scabby potato field in different regions of Egypt. The taxonomy of the organism was determined by morphology, biochemistry, and physiological characteristics. Two bacterial strains (actinomycetes) which were isolated from soil sample and showed antagonistic activity towards potato scab were identified using specific polymerase chain reaction (PCR) of 16S rDNA gene. The 16SrDNA sequence analysis showed that the 1st strain belongs to the genus *Streptomyces*, with closest similarity to *Streptomyces avermitilis* MA-4680 (100% similarity). Sequence similarities between the 1st strain and other *Streptomyces* species in the same subclade ranged from 98% (with *Streptomyces griseus* NBRC 13350 and *Streptomyces* sp. Wigar10). Key Phenotypic characteristics as well as chemotaxonomic features of the actinomycetes were congruent with the description of the genus *Streptomyces*. On the basis of phenotypic and phylogenetic analyses. The 2nd strain identified as *Actinomyces odontolyticus* ATCC 17982(100% similarity). Sequence similarities between the 2nd strain and other Actinomycetes species was 98% (with *Actinomyces odontolyticus* C 505). Both identified strains showed a high level of antibiosis against pathogenic organism (*Streptomyces scabies*) and the completely potato controlled the disease as no sign of disease symptoms were shown on the tested varieties (Cara and Diamond)

Key words: Actinomycetes, biological control, *Streptomyces scabies*.

INTRODUCTION

Potato (*Solanum tuberosum* L.) is one of the most important vegetable crops in Egypt, for both local consumption and exportation. It is widely cultivated and could contribute to reduce worldwide food shortages (Han *et al.*, 2005). One of the most important potato diseases is common scab caused by *Streptomyces scabies*. It is indigenous in all potato growing areas in the world (Loria *et al.*, 2006; Wanner, 2004). Several species of *Streptomyces* can cause PCS but *Streptomyces scabies* is considered to be predominant (Lambert and Loria, 1989). It can infect number of root crops, including radish, parsnip beet and carrot (Wanner, 2004). The disease has little impact on total potato yield but spoils the appearance, quality, and market ability of the tubers (Johnson and Powelson, 2008). Several methods have been used to control potato

scab, including: 1) Planting resistant varieties and 2) agriculture practices such as excess irrigation during tuber formation (Lapwood and Adams, 1975). However all these methods are less effective and durable with seldom preventing disease from occurring but generally reducing its extent of severity 3- Chemical control, such as (3,5-Dtelon) and (polyram and mancozeb). However these chemical increase phytotoxicity to freshly cut tuber and decrease in the tuber size and yield. Therefore, we need more research into nonchemical methods of potato production seems to be justified than other. Biological control is suitable alternative of chemical control.

Microorganisms are virtually unlimited sources of novel compounds with many medicinal and agricultural applications. Actinomycetes, among them, hold a

prominent position due to their ability to produce numerous different metabolites such as antibiotics, enzymes and inhibitors (Xu *et al.*, 2005). Further; the discovery of novel antibiotic and nonantibiotic lead compounds through microbial secondary metabolite screening is becoming increasingly important. In recent years; there has been an increasing interest in discovering new agricultural antibiotics for the protection of our living environments. The genus *Streptomyces* is the largest producer of bioactive compounds (Chun *et al.*, 1997; Labeda *et al.*, 1997).

Actinomycetes are gram-positive, aerobic, high GC-content and 0.5–1.0 µm in size. They are filamentous, sporulating colonies and recognized as a transition group between primitive bacteria and fungi (Lo *et al.*, 2002). Among the actinomycetes groups, *Streptomyces* are the most popular and found worldwide in soil, and important in soil ecology. They belong to the order Actinomycetales. *Streptomyces* are metabolically diverse and can utilize almost anything as carbon source due to its ability to produce extracellular hydrolytic enzymes, including sugars, alcohols, amino acids, organic acids, aromatic compounds and other complex substrates such as cellulose, mannan and xylan. They are also well known for their abilities to produce antibiotics and other secondary metabolites (Willey *et al.*, 2008). Thus, these microorganisms have been implicated in the antagonism of a wide variety of plantpathogenic bacteria, fungi and nematodes for their potential use as biological disease control agents (Sahilah *et al.*, 2010).

In the context of the information provided above, the objectives of this study have been, to evaluate the potential of these two antibiotic-producing suppressive strains of Actinomycetes to control potato scab disease.

MATERIALS AND METHODS

Isolation of *Streptomyces scabies* (pathogen). Samples of soils and tubers of potato cultivars with scab symptoms were collected from fields of different locations of

five Egyptian governorates (Kafr El-Sheikh, El-Gharbiya, El-Minufya, El-Dakahlia and El-Nobaria). Isolation of actinomycetes from the scab lesions was carried out using the methods described by Lawrence (1956). Cultural, morphological microscopical and pathological properties were considered to identify the isolated pathogens according to Burgess *et al.* (1994).

Morphological and physiological characterization of *Streptomyces scabies* (pathogen). The morphology of the sporophores was examined microscopically, and the color of spores and colonies were observed on oatmeal agar (OMA) after 14 days of incubation at 28°C (Holt, *et al.*, 1994). Production of soluble pigments were observed after 4 days of incubation at 28°C. The ability to utilize the International Streptomyces Project (ISP) sugars was tested on the standard basal medium (Shirling and Gottlieb, 1966).

Pathogenicity tests:

Pathogenicity tests were performed on potato cultivars using the methods described by Labruyere (1971). Thirty four strains were isolated and tested for pathogenicity. Inocula were prepared by growing each tested strain for two weeks at 30°C in 50ml tubes containing sterilized vermiculite saturated with a modified Say – solution composed of 20gm of Sucrose, 1.2gm of L-asparagine, 0.6gm of K₂HPO₄ and 10 gm of yeast extract per liter of water. Healthy potato cultivars Cara and Diamond were individually planted in 35cm diameter pots containing sterile clay soil mixed with 20ml of the inoculum. Plants were arranged in a growth chamber as a randomized complete block with three replicates. Uninoculated controls were included in the tests. Potatoes were harvested after three months and tubers were examined for common scab symptoms.

Isolation and screening of antagonists.

Antagonists were isolated from soil rhizosphere samples of healthy potato plants producing areas at some governorates of Egypt. The used bioagents

Identification and characterization of actinomycetes for biological

were isolated on selected media according to the methods recommended by Anonymous (1984), Burgess *et al.* (1994) and Turner *et al.* (1998).

Biological control of the tested microorganisms on potato scab pathogen.

In vitro

The antagonistic activity was estimated by disc diffusion method (Barakate *et al.*, 2002). A disc of 5mm in diameter from every microorganism (48 hours old culture) was placed on the surface of OMA plates seeded with potato scab pathogen. The plates were incubated at 28 °C for 24 h. The inhibition zone around the discs indicated the antagonistic interaction.

In vivo

Pots of 30 cm in diameter were filled with nonsterilized clay-loam soil at the rate of 8 Kg per pot. Soil infestation was carried out using the pathogen *Streptomyces scabies* (control) and antagonistic microorganisms namely (*Actinomycetes odontolyticus*, *Streptomyces avermitilis*, *Streptomyces griseus*, *Bacillus subtilis*, *Bacillus thuringiensis*, *Trichoderma hamatum* and *Trichoderma koningii*) according to (Michel and Mew, 1998). A tuber of either potato cultivars (Cara and Diamond) was planted per pot (a replicate). The same eight treatments were repeated with the other cultivar of potato. Each treatment was represented by six replicates. At harvest the average weight of tuber and percentage of diseased tubers were recorded.

Statistical analysis.

The obtained data were statistically analysed according to the method of Gomez and Gomez, (1984).

Molecular characterization of *Streptomyces avermitilis* and *Actinomycetes odontolyticus*

- Genomic DNA extraction from actinomycetes and *Streptomyces*. DNA extraction was carried out following the CTAB method according to (Azadeh and Meon, 2009). Ten colonies of bacteria was

inoculated into 10 mL nutrient broth and incubated at 28±2°C overnight. One mL of the overnight culture was transferred into a 1.5 mL eppendorf tube and centrifuge for 30 sec at 13,000 rpm. The bacteria cells were collected by discarding the supernatant and resuspended in 567 µL TE buffer (10 mM Tris-HCL, pH 7.4, 1 mM EDTA and 1 L distilled water), mixing well by vortexing. Then 30 µL of 10 % SDS, 3 µL of NAOAC (sodium acetate) pH 5.2, 100 µL of 5 M NaCl and 80 µL CTAB-NaCl were added to a total volume of 780 µL and mixed well before incubating for 10 min in water bath at 65°C. An equal volume (780 µL) of chloroform/isoamyl alcohol (24:1) was added to the mixture and centrifuged at 13000 rpm for 5 min to separate the phases. The clear supernatant was transferred into a new eppendorf tube and the aqueous DNA layer was again extracted using phenol/chloroform/isoamyl alcohol (25:24:1). This step was repeated 3 times and the supernatant pooled. The clear supernatant was transferred into new eppendorf tube and 400 µL of isopropanol was added to precipitate the nucleic acid. Finally the DNA was washed with 200 µL of 75% of cooled ethanol and dried at room temperature (28±2°C) before dissolving in 100 µL sterile distilled water and kept at -20°C for further analysis. Electrophoresis was run for identifying the nucleic acids after DNA extraction in 1% of agarose gel and 1% of TBE (Tris base, boric acid, 0.5 M EDTA solution, 1L ddH₂O at pH 8.0). The products were mixed with loading dye buffer (MBI Fermentas) in 5:1 ratio and subjected to electrophoresis at 70 volts for 1 h and 45 min. DNA ladder 100 bp (MBI Fermentas) was used as marker. The gel was stained in ethidium bromide solution and the bands visualized and photographed using Sony digitalcamera.

Primer design and PCR amplification. Oligonucleotide primer for 16S rDNA gene was 16S-1f (5'- GCTAGTTGGTGGGGTAA-3', 17 mer) and 16S-2r (5'- GCCATCTCAGTTCGGATTG-3'; 18 mer) were designed on the basis of the sequence of *E. coli* 16 S gene (corresponding to positions 247 to 263 and 1291 to 1309; *E coli* numbering system) (Wilems and Collins, 1993).

Oligonucleotide primer for Glyphosate tolerant gene was AroA f (5'-GCTCTAGAAGTGTGGAACAATATG-3'; 27 mer) and AroA r (5'-TTACTCGAGTGAGAATTAATTGATGG-3'; 33 mer) (Sun *et al.*, 2005). Amplification reaction for bacteria was performed in 25 µL of total volume containing 2 µL of DNA as a template, 2.5 µL of 10 PCR buffer (Fermentas), 1.5 µL of 25 mM MgCl₂ (Fermentas), 0.2 µL of 10 mM dNTPs, 0.1 µL primer oligonucleotides, 0.1 µL Taq polymerase (Fermentas) and 18.5 µL of sterilized distilled water. The amplification was performed in a Thermal Cycler (Biometra®, T3 thermocycler) (Syngene, UK) programmed for pre-denaturing of 3 min at 94°C, 30 cycles of 1min at 94°C, 1min at 58°C and 2 min at 72°C. After a final extension of 7 min at 72°C, the samples were cooled to 4°C.

Nucleotide sequence analysis: The PCR products were purified using a commercial kit (QIA Quick PCR purification kit (Qiagen, Valencia, CA), according to manufacturer's instruction. After purification, the PCR products were sent for sequencing services at Sigma Co. Germany. The 16 S gene sequences were aligned using BioEdit software version 7.0.8 (<http://www.mbio.ncsu.edu/bioEdit/bioEdit>) and searched for sequence similarity to other sequences which are available in the NCBI database at <http://www.ncbi.nih.gov> using Basic Local Alignment Search Tool (BLAST) algorithm. Multiple sequence alignments were performed on the selected closely related sequence accessions available using bioedit software (<http://bioedit.edu/>).

Phylogenetic analysis:

Phylogenetic analysis was done based on the nucleotide sequences of 16 S gene using draw tree software provided by the Biology Workbench Program (<http://workbench.sdsc.edu/>).

Number of Data base JYZVMFRR015, K12FNS3P01S

RESULTS AND DISCUSSION

Isolation of *Streptomyces scabies* (pathogen).

Streptomyces scabies Waksman & Henrici occurs worldwide and causes common scab on potato tubers (Hooker, 1981). The type strain of *S. scabies* (ATCC 49173) (Lambert and Loria, 1989) is characterized by gray spores born in spiral chains (sporophores), melanin pigment production on tyrosin-containing medium (peptone iron agar), and utilization of all the diagnostic sugars recommended by the International Streptomyces Project (ISP) (Shirling and Gottlieb, 1966) for identification of *Streptomyces* spp. *Streptomyces scabies* does not grow at pH 4.5 (Lambert and Loria, 1989). Subsequently, common scab is less severe in acidic soils and does not usually occur in soils of pH < 5.2. Severity of common scab can also be reduced by maintaining high soil moisture with irrigation and by growing resistant cultivars (Lindholm *et al.*, 1997).

Description of Streptomyces isolated from scabby potato plants. *Streptomyces* spp. were isolated from scabby potato plants collected from fields of different locations from many governorates, Egypt (Table 1). The original scabby tubers showed a range of symptoms from superficial small discrete lesions covering parts of a tuber to large, deep, coalescing raised or pitted lesions covering most or all of a tuber. They also varied in their pathogenicity on radish and on potato cultivars.

Morphological and physiological characterization of *Streptomyces scabies* (pathogen).

Morphological observation of the 7–15 days old culture of strain isolated from potato scab is revealed that both aerial and vegetative hyphae were abundant, well developed. Long spore chains were developed on the aerial mycelium. The aerial mycelium was observed after the 15th day of incubation in all tested media. Cultural characteristics of the strain are shown in (Table 2). Aerial mycelium of this strain was abundant, well-developed and varied from brown to white on different

Identification and characterization of actinomycetes for biological

tested media. The substrate hyphae varied from pale-yellow to brown or white. Yellow diffusible pigments were produced on Yeast extract–malt extract agar media, and melanin

was produced on Peptone- Yeast extract-iron agar.

Physiological and biochemical characteristics of the same strain isolated from potato scab are indicated in Table 3.

Table 1: Number of the obtained *Streptomyces scabies* isolates from different governorates and their pathogenicity reaction:

Region	Isolate source	Scab type	No. of isolates	Pathogenicity	
				Potato tubers	Radish seedling
Kafr El-Sheikh	Tuber	Common	6	+	+
Kafr El-Sheikh	Soil	-	3	+	+
El-Gharbiya	Tuber	Common	8	+	+
El-Gharbiya	Soil	-	4	+	-
El-Minufya	Tuber	Superficial	3	+	-
El-Minufya	Soil	-	1	+	+
El-Dakahlia	Tuber	Common	7	+	+
El-Dakahlia	Soil	-	0	+	+
El-Nobaria	Tuber	Superficial	2	+	+
El-Nobaria	Soil	-	0	+	-

+ Positive reaction

- Negative reaction

Table 2: Cultural characteristics of strain *Streptomyces scabies*.

Medium	Color of Aerial mycelium	Color of Substrate mycelium	Production of soluble pigment
Yeast extract–malt extract agar (ISP medium 2)	Brown	Yellow	Yellow
Oatmeal agar (ISP medium 3)	White	White	None
Peptone- Yeast extract- iron agar (ISP medium 6)	Brown	Brown	Melanin
Tyrosine agar (ISP medium 7)	Yellowish White	White	None
Nutrient agar	White	White	None
Czapek's agar	White	White	None

ISP, International Streptomyces Project Shirling and Gottlieb (1966).

Table 3 : Morphological and Physiological characterization of *Streptomyces scabies*.

Characteristic	Result
Shape of cell	Spiral hyphae
Size of cell	Long
Arial mycelium colour	Grey to brown
Colour of spores	Grey
Gram's staining	+ve
Pigment production agar	brown
Optimum temp.	28-30o
Utilization of sugars:	
Mannitol	++
Fructose	+
Sucrose	++
Arabinose	+
Glucose	++
Fructose	+
Raffinose	+
Rhamnose	+
Starch hydrolysis	-
Gelatin liquefaction	+
Indol formation	-
Catalase activity	+
V.P. Test	+
Methyl Red (MR)	-
Production of H ₂ S	-
Nitrogen reaction	-
Litmus milk	+

+ Positive Utilization
++ strong positive Utilization
- negative reaction

These results are in harmony with those obtained by Lindholm *et al.*, 1997 which showed that *S. scabies* (type strain ATCC 49173) is characterized by spiral

Identification and characterization of actinomycetes for biological

sporophores and gray spores, produces of melanin pigment on peptone-iron agar, utilizing of all ISP sugars, and pathogenic on potato. Also color of colonies and production of diffusible pigments are criteria for species identification.

Pathogenicity test : Results obtained clear that potato common scabe symptoms varied according to *Streptomyces scabies*.

Isolation and screening of the antagonists: Three Actinomycetes (*Actinomycetes odontolyticus*, *Streptomyces avermitilis* and *Streptomyces griseus*), 2 bacteria (*Bacillus subtilis* and *Bacillus thuringiensis*) and 2 fungi (*Trichoderma hamatum* and *Trichoderma koningii*) were isolated. The diameter of inhibition zone is considered as criterion of antagonistic efficiency of biocontial agent.

Biological control of the tested microorganisms on potato scab pathogen.

***In vitro* studies:**

The obtained isolates (3 streptomyces and 2 bacteria and 2 fungi) were screened against the plant pathogen *Streptomyces scabies* and showed antagonism activity against it as given in Table (4).

In vivo

At harvest, the average weight of tuber and percentage of diseased tuber were determined. The two strains that showed most clearing zone towards *S. scabies* (*Streptomyces avermitilis* and *Actinomycetes odontolyticus*) resulted the highest average weight of tuber and also decreased the percentage of disease incidence on tubers (a high biological control of disease) as shown in Table (5).

Table 4: Antagonistic efficiency between the isolated microorganisms and *S. scabies*:

Antagonism	Inhibition zone in mm
<i>Actinomycetes odontolyticus</i>	48.9
<i>Streptomyces avermitilis</i>	51.6
<i>Streptomyces griseus</i>	32.0
<i>Bacillus subtilis</i>	24.9
<i>Bacillus thuringiensis</i>	24.3
<i>Trichoderma hamatum</i>	19.6
<i>Trichoderma koningii</i>	22.9

Table 5: the average weight of tuber and percentage of diseased tubers:

Organisms	Average weight of tuber		% of diseased tubers	
	Diamond	Cara	Diamond	Cara
<i>Streptomyces scabies</i>	76.67AB	53.57B	96.89A	94.14A
<i>Actinomycetes odontolyticus</i>	79.32 A	58.38 A	0.00 F	0.00 E
<i>Streptomyces avermitilis</i>	78.08 A	53.45 B	0.00 F	0.00 E
<i>Bacillus subtilis</i>	66.79 C	50.35 C	2.147 E	0.00 E
<i>Bacillus thuringiensis</i>	64.28 CD	48.67 D	9.443 D	5.803 D
<i>Streptomyces griseus</i>	74.01 B	53.57 B	31.51 B	22.13 B
<i>Trichoderma koningii</i>	62.04 D	40.15 E	16.27 C	15.16 C
<i>Trichoderma hamatum</i>	70.00 A	50.01 B	50.12 A	23.21 B
L.S.D	3.604 at α 0.05		0.689 at α 0.05	

It is worthy to note that the strains of *Streptomyces avermitilis* and *Actinomycetes odontolyticus* showed high protective of antibiosis against Pathogen are subjected to further identification (morphological cultural and biochemical characteristic) (Tables 6 and 7).

Morphological observation of the 7–15 days old culture of strains isolated and showed antagonism against potato scab is revealed that both aerial and vegetative hyphae were well developed. Long spore chains were born on the aerial mycelium. The aerial mycelium was observed after the 15th day of incubation on all tested media. Cultural characteristics of the two strains are shown in Table 5. Aerial mycelium of strain 1 was abundant, well-developed and varied from white to brown on different tested media. The substrate hyphae varied from white to brown. Aerial mycelium of strain 2 was well-developed and varied from white to orange- brown on the tested media. The substrate hyphae varied from white to

brown. The diffusible pigment melanin was observed on ISP-6 medium.

These results also were consistent with results obtained for 16SrDNA gene sequences DNA, indicated that one of the two spp. is belongs to actinomycetes and the other is represented as *Streptomyces* sp.

Identification of the two spp. using 16SrDNA partial sequences DNA.

Two actinomycetes were examined for the specific amplification of 16S rDNA gene sequences (Azadeh and Meon, 2009). As indicated in Table 4, the most active microorganisms against potato scab were strains 1 and 2. They were identified as *Streptomyces* species with 100% sequence similarity. Strain 1 and strain 2 were identified up to species level, namely *Streptomyces avermitilis* MA-4680 and *Actinomycetes odontolyticus* ATCC 17982 respectively.

Analysis of the 16SrDNA from *Streptomyces* species showed that the 1st

Table 6: Cultural characteristics of the two antagonistic strains to *Streptomyces scabies*

Medium	Color of Aerial mycelium		Color of Substrate mycelium		Production of soluble pigment	
	Strain 1	Strain 2	Strain 1	Strain 2	Strain 1	Strain 2
Yeast extract–malt extract agar (ISP medium 2)	Brown	Orange-brown	Yellow	Brown	Yellow	Yellow
Oatmeal agar (ISP medium 3)	White to yellow	White to yellow	White	White	None	None
Peptone- Yeast extract-iron agar (ISP medium 6)	Brown	Brown	Brown	Brown	Melanin	Melanin
Tyrosine agar (ISP medium 7)	Yellowish White	Yellow	White	White	None	None
Nutrient agar	White	Brown	White	Yellow	None	None
Czapek's agar	White to yellow	Brown	White	Yellow	None	None

ISP, International Streptomyces Project Shirling and Gottlieb (1966).
Strain 1: *Streptomyces avermitilis* strain 2: *Actinomycetes odontolyticus*

Identification and characterization of actinomycetes for biological

Table 7: Morphological and Physiological characterization of the two antagonistic organisms.

Characteristic	Strain 1	Strain 2
Shape of cells	Hyphae	Hyphae
Size	Long	Long
Gram's staining	+	+
Sporulation	+	+
Motility	-	-
Growth of KBA Medium	-	-
Optimum temp.	28-30°	30°
Utilization of sugar:		
Mannitol	AG	A
Fructose	-	A
Sucrose	AG	AG
Arabinose	-	A
Glucose	A	A
Galactose	A	A
Lactose	-	-
Maltose	A	AG
Dextrose	AG	AG
Glycerol	A	A
Menthol	-	A
Raffinose	-	-
Starch hydrolysis	-	-
Gelatin liquefaction	+	+
Indole formation	-	-
Catalase activity	+	+
Lipolytic activity	+	-
V.P. Test	+	+
Methyl Red (MR)	-	-
Production of H ₂ S	-	-
Nitrate reduction	-	+
Milk coagulation	+	+

(+) Positive reaction (-) Negative reaction G: Gas A: Acid
 Strain 1: *Streptomyces avermitilis* strain 2: *Actinomycetes odontolyticus*

Strain was grouped into a branch with *Streptomyces coelicolor* A3 (similarity value of 97%; the closest neighbors). The almost complete 16S rRNA gene sequence of 1st strain was determined in this study and has

been deposited in the GenBank database. This sequence was subjected to similarity searches against public databases to infer a possible phylogenetic relationship of this strain. This analysis revealed that this strain

was a member of the genus *Streptomyces*. A neighbor-joining tree (Saitou and Nei, 1987) (Figure 1) based on 16S rDNA gene sequences were constructed to show relationships between the strain and some other related *Streptomyces* species. Bootstrap analysis was used to evaluate the tree topology of the neighbor-joining data by performing 1000 resampling (Felsenstein, 1985). Thus, based on the results of the above phenotypic and genotypic analyses, strain 1 should represent a species of the genus *Streptomyces*, for which we propose the name *Streptomyces avermitilis* MA-4680.

Description of *Streptomyces avermitilis* MA-4680

Gram positive organism both vegetative and aerial hyphae were abundant and well-developed, spore forming. Aerial mycelium varied from white to brown. The substrate hyphae from white to brown. The diffusible pigment melanin was observed on ISP-6 medium. Gelatin liquefaction, Catalase activity, Lipolytic activity, V.P. test and milk coagulation were positive while starch hydrolysis, H₂S and indole production, nitrate reduction were negative. Glucose, mannitol, sucrose, galactose, maltose, dextrose and glycerol are utilized but not

fructose, lactose, arabinose, raffinose and mannitol.

The analysis of the 16SrDNA for 2nd strain showed that *Streptomyces* species could be grouped into a branch with *Actinomyces odontolyticus* C 505(similarity value of 98%; the closest neighbors).The almost complete 16S rRNA gene sequence of 2nd strain was determined in this study and has been deposited in the GenBank database. This sequence was subjected to similarity searches against public databases to infer possible phylogenetic relationships of this strain. This analysis revealed that this strain was a member of the genus actinomycetes. A neighbor-joining tree (Saitou and Nei, 1987) as shown in Figure (2) based on 16S rDNA gene sequences were constructed to show relationships between strain 2 and some other related actinomycetes species. Bootstrap analysis was used to evaluate the tree topology of the neighbor-joining data by performing 1000 resamplings (Felsenstein, 1985). Thus, based on the results of the above phenotypic and genotypic analyses, strain 2 should represent a species of the genus actinomycetes, for which we propose the name *Actinomyces odontolyticus* ATCC 17982.




Figure 1. Phylogenetic dendrogram obtained by distance matrix analysis of 16S rDNA sequences, showing the position of strain *Streptomyces avermitilis* MA-4680 among phylogenetic neighbors. Numbers on branch nodes are bootstrap values (1000 resamplings).




Figure 2. Phylogenetic dendrogram obtained by distance matrix analysis of 16S rDNA sequences, showing the position of strain *Actinomyces odontolyticus* ATCC 17982 among phylogenetic neighbors. Numbers on branch nodes are bootstrap values (1000 resamplings).

Description of *Actinomyces odontolyticus* ATCC 17982

Gram positive organism both vegetative and aerial hyphae were abundant and well-developed; spore forming. Aerial mycelium of strain 2 varied from white to orange-brown on the tested media and the substrate hyphae from white to brown. The diffusible pigment melanin was observed on ISP-6 medium. Gelatine liquification, Catalase activity, nitrate reduction, V.P. test and milk coagulation were positive while starch hydrolysis, H₂S and indole production were negative. Glucose; mannitol, fructose, sucrose, arabinose, galactose, maltose, dextrose and glycerol are utilized but not lactose and raffinose.

Results of the present study provide sufficient evidence to recommended use of *Actinomyces odontolyticus* and *Streptomyces overmitils* as successful biocontrol agents against *Streptomyces scabies* causing common scab disease on both tested varieties of potatoes (Diamond and Cara).

REFERENCES

- Anonymous (1984). Bergey's Manual of Systematic Bacteriology. Ordinary gram positive bacteria. Vol. 1. Krieg, N.R. (ed.). Ordinary gram negative bacteria. Vol. 2, Sneath, P.H.A. (ed.). Williams and Wilkins, Baltimore, USA.
- Azadeh, B. and S. Meon (2009). Molecular Characterization of *Pseudomonas aeruginosa* UPM P3 from Oil Palm Rhizosphere. *Amer. J. Appl. Sci.* 6 (11): 1915-1919.
- Barakate, M., Y. Ouhdouch, K. Oufdou and C. Beaulieu (2002). Characterization of rhizospheric soil streptomycetes from Moroccan habitats and their antimicrobial activities. *World Journal of Microbiology and Biotechnology* 18: 49-54.
- Burgess, L.W., B.A. Summerell, S.G. Bullock and K.P.D. Backhouse (1994). Laboratory Manual for Fusarium Research. 3rd Ed. University of Sydney. 133pp.
- Chun, J., H.D. Youn, Y.I. Yim, H. Lee, M.Y. Kim, Y.C. Hah and S.O. Kang (1997). *Streptomyces seoulensis* sp. nov. *Int. J. Syst. Bacteriol.* 47: 492-498.

- Felsenstein, J. (1985). Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39: 783–791.
- Gomez, K.A. and A.A. Gomez (1984). *Statistical Procedures for Agricultural Research*. 2nd Ed. John Wiley and Sons, pp. 229-308
- Han, J.S., J.H. Cheng, T.M. Yoon, J. Song, A. Rajkarnikar, W.G. Kim, I. D. Yoo, Y.Y. Yang and J.W. Suh (2005). Biological control agent of common scab disease by antagonistic strain *Bacillus* sp. sunhua. *J. Appl. Microbiol.*, 99: 213-221.
- Holt, J. G., N. R. Krieg, P. H. A. Sneath, J. T. Staley and S. T. Williams (1994). *Bergey's Manual of Determinative Bacteriology*, 9th ed. Williams and Wilkins, Baltimore, MD.
- Hooker, W. J. (1981). Common scab. Pages 33-34 in: *Compendium of Potato Diseases*. American Phytopathological Society, St. Paul, MN.
- Johnson, D. A. and M. L. Powelson (2008). *Potato Health Management*. American Phytopathological Society Press, St. Paul, MN.
- Labeda, D.P., M.P. Lechevalier and R.T. Testa (1997). *Streptomyces stramineus* sp. nov., a new species of the verticillate streptomycetes. *Int. J. Syst. Bacteriol.* 47: 747–753.
- Labruyere, R.E. (1971). Common scab and its control in seed, potato crops. Center for Agricultural Publishing and Documentation, Wageningen, Netherlands.
- Lambert, D. H. and R. Loria (1989). *Streptomyces scabies* sp. nov. nom. Rev. *Int. J. Syst. Bacteriol.* 39:387-392.
- Lapwood, D. H. and M. J. Adams (1975). Mechanisms of control of common scab by irrigation. P123-129 in: *Biology and Control of soil borne Pathogens*. G. W. Bruehl, ed. The American Phytopathological Society, St. Paul, MN.
- Lawrence, C. H. (1956). A method of isolating Actinomycetes from scabby potato tissue and soil with minimal contamination. *Can. J. Bot.* 34:44-47.
- Lindholm, P., H. Kortemaa, M. Kokkola, K. Haahtela, M. Salkinoja-Salonen and J. P. T. Valkonen (1997). *Streptomyces* spp. isolated from potato scab lesions under Nordic conditions in Finland. *Plant Dis.* 81:1317-1322.
- Lo, C.W., N.S. Lai, H-Y. Cheah, N.K.I. Wong and C.C. Ho (2002). Actinomycetes isolated from soil samples from the Crocker Range Sabah.
- ASEAN Review of Biodiversity and Environmental Conservation* (ARBEC).
- Loria, R., J. Kers and M. Joshi (2006). Evolution of plant pathogenicity in *Streptomyces*. *Annu. Rev. Phytopathol.* 44:469-487.
- Michel, V.V. and T.W. Mew (1998). Effect of soil amendment on the survival of *Rastonia solanacearum* in different soil. *Phytopathology* 88: 300-305.
- Sahilah, A.M., S.Y. Tang, M.N. Zaimawati, H. Rosnah, M.S. UmiKalsum and R. Son (2010). Identification and characterization of actinomycetes for biological control of bacterial wilt of *Ralstonia solanacearum* isolated from tomato. *J. Trop. Agric. and Fd. Sc.* 38 (1) 103-114.
- Saitou, N. and M. Nei (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* 4: 406–425.
- Shirling, E. B. and D. Gottlieb (1966). Methods for characterization of *Streptomyces* species. *Int. J. Syst. Bacteriol.* 16:313-340.
- Sun, Y.C., Y.C. Chen, Z. X. Tian, F. M. Li, X. Y. Wang, J. Zhang, Z. L. Xiao, M. Lin, N. Gilmartin, D. N. Dowling and Y. P. Wang (2005). Novel AroA with high tolerance to Glyphosate, encoded by a gene of *Pseudomonas putida* 4G-1 isolated from an extremely polluted environment in China. *Appl. Environ. Microbiol.* 71: 4771–4776.
- Turner, D., W. Kovacs, K. Kuhls, E. Lieckfeldt, B. Peter, I. Arisan Atac, J. Strauss, G.J. Samuels, T. Borner and C.P. Kubicek (1998). Biogeography and phenotypic variation in *Trichoderma* sect. *longibraclatum* and associated *Hypocrea* species. *Mycol. Res.*, 101:449-549.
- Wanner, L. A. (2004). Field isolates of *Streptomyces* differ in pathogenicity and virulence on radish. *Plant Dis.* 88:785-796.

Wilems A. and M. D. Collins (1993). Phylogenetic analysis of Rhizobia and Agrobacteria based on 16s rRNA gene sequences. Int. J. Syst. Microbiol. 43: 305-313.
Willey, J.M., L.M. Sherwood and C.J. Woolverton (2008). *Prescott, Harley and*

Klein's Microbiology, 7th Edn. USA: McGraw Hill Higher Education.
Xu, L., Y. Jiang, W. Li, M. Wen, M. Li and C. Jiang (2005). *Streptomyces roseoalbus* sp. nov., an actinomycete isolated from soil in Yunnan, China. *Antonie van Leeuwenhoek*, 87:189–194.

المقاومة الحيوية لمرض الجرب في البطاطس بواسطة عزل وتعريف نوعين من الاكتينوميستات

هند عبد الحميد حميدو (1) وعبير حمدي مخلوف (2)

(1) كلية التربية بالعريش - جامعة قناة السويس

(2) كلية الزراعة - جامعة المنوفية.

المُلخَص العربي:

يعتبر مرض الجرب في البطاطس والمتسبب من بكتيريا الاستريتومايسس اسكابس من أخطر الأمراض التي تصيب أحد المحاصيل الاقتصادية بمصر. تم عزل المسبب للمرض من درنات البطاطس المصابة وتعريفه كنوع جديد من مسببات الجرب في مصر من الاستريتومايسس وتم دراسة الصفات التقسيمية والظاهرية وإجراء التجارب الفسيولوجية والبيوكيميائية لتعريفه والتي أظهرت أنه من نوع الاستريتومايسس اسكابس. تم عزل عزلات مضادة فطرية وبكتيرية من تربة الريزوسفير لنباتات بطاطس سليمة وتم اختبار قدرتها التضادية للمسبب لمرض الجرب وأوضحت النتائج أن أكثر العزلات مقاومة لهذا المرض عزلتين من جنس الاكتينومييسس. وقد تم تعريفهما بواسطة التجارب الفسيولوجية والبيوكيميائية ، كذلك الصفات الظاهرية وللتأكيد تم تعريفها جينيا بواسطة تفاعل البلمرة المتسلسلة وقد عرف الكائن الأول بأنه (*Streptomyces avermitilis* MA-4680) والثاني (*Actinomyces odontolyticus* ATCC 17982) وأوضحت نتائج المقاومة الحيوية للمسبب لمرض الجرب تحت ظروف العدوى في الصوبة الزجاجية بواسطة العزلات المضادة والسابقة الذكر قد قللت المرض بصورة معنوية وأدت الي زيادة ملحوظة في وزن درنات البطاطس المختبرة لنوعي (الكارا والدياموند).