

## **EFFECT OF ADDITION DIFFERENT LEVELS OF DRIED ONION AND GARLIC ON DIGESTIBILITY, CAECOTROPHY AND CAECUM ACTIVITY OF RABBITS**

El-Sayaad, G. A. E.<sup>1</sup> ; A. A. Gabr<sup>2</sup>; and W. A. E. Ibrahim<sup>2</sup>

<sup>1</sup> Fac. of Agric., Benha Univ.

<sup>2</sup> Fac. of Agric., Mansoura Univ.

### **ABSTRACT**

A total number of 42 male New Zealand white (NZW) rabbits at the age of 14 weeks were randomly divided into seven experimental groups (6 male for each) to study the effect of adding dried onion ( DO) and garlic (DG) on nutrients digestibility , caecotrophy and caecum activity. Rabbits were fed a basal diet supplemented with different levels of DO and DG as follows: 1) control group (CG), 2) CG+1 % DO, 3) CG+1.5 % DO, 4) CG+1% DG, 5) CG+1.5 % DG, 6) CG +0.5 % DO+0.5% DG, 7) CG +0.75% DO +0.75 % DG, respectively. All diets were formulated to be iso-nitrogenous and iso-caloric. The obtained data showed that dry matter intake did not significantly differ among dietary treatments as well as it was similar for collared rabbits compared with those uncollared. The CP digestibility coefficient was significantly ( $p<0.05$ ) higher for rabbits fed diets included 1.5 % DG, 1.5 % DO, 0.5 % DO+ 0.5% DG and 1 % DG than those fed either control diet or other groups. Rabbits fed 0.5% Do + 0.5% DG diet recorded significantly higher CF digestibility ( $P<0.05$ ) than the other treated groups. The nutritive values expressed as TDN%, DE and ME Kcal/Kg DM of diet supplemented with 0.5% DO +0.5% DG were significantly ( $p<0.05$ ) higher than those fed 1% Do. The DCP% for 1.5% DO, 1% DG, 1.5%DG and 0.5% DO+0.5% DG diets was significantly ( $p<0.05$ ) higher than 1% DO, 0.75% DO+0.75%DG and control groups. The digestibility coefficients of all nutrients as well as nutritive values of tested diets were similar between uncollared and collared groups .Treatment groups showed significant ( $p<0.05$ ) increase in Bacterial total count compared with the control group.

**Keywords:** rabbits, onion, garlic, digestibility, nutritive value, caecotrophy, caecum activity.

### **INTRODUCTION**

Modern animal production requires the use of safe and effective additives to stimulate feed consumption and destroy harmful microorganisms of the diet, in addition to be used as rumen manipulators to increase animal productivity (Ahmed *et al*, 2009). Several attempts were conducted to use natural materials such as medicinal plants as feed additives could be widely accepted (line-Eric *et al*; 1998 and Aboul-fotouh *et al*; 2000). The World Health Organization (WHO) encourages using medicinal herbs and plants to substitute or minimize the use of chemicals through the global trend to go back to the nature (El-Ashry *et al* 2006). Many investigators reported that garlic and onion are highly inhibitory to E.coli and to other bacteria and fungi, e.g. antibacterial and antifungal (Wager *et al*; 1994 and Kumar and Berwal 1999). Nutritional activities of garlic and onion have been widely studied. The active inhibitory agents of garlic are allicin and/or daily thiosolphinic acid

allicin which is enzymatically released from precursor form when the garlic and onion bulbs are crushed (Saleem and Al-Delaimy 1982). With rabbit (El-elamei 2001; Helal and Mohamed 2001; El-Hindawy *et al* 2003 and Abdel-Azeem and Abdel-Reheem 2006) and crossbred heifers and growing buffalo calves (Gupta *et al* 2005 and Aiad *et al* 2008). Onion and garlic supplementation showed an improvement in animal performance, gross activity of caecum or rumen microflora. However, such effects of these additives could be differ according to many variables e.g. type and level of these additive, animal species, plane of nutrition ,management conditions. Therefore, the present study was designed to evaluate the effects of onion and garlic addition as natural safe feed additives in rabbit diets on digestibility, caecotrophy status and caecum activity.

## **MATERIALS AND METHODS**

The present study was carried out at the Rabbit's Farm, Department of Animal Production, Faculty of Agriculture, Moshtohor, Banha University, Egypt. The experiment was done during the period from May to July 2009.

### **Experimental diets**

The composition and calculated chemical analysis of the experimental diets are presented in Table (1).

Diets were formulated by mixing dried onion (DO), dried garlic (DG) and the mixture of both at levels 1, 1.5%, (0.5% DO + 0.5 %DG) and (0.75 % DO + 0.75 % DG), respectively. Seven experimental diets were formulated to be iso-caloric and iso-nitrogenous. All diets and supplements covered the daily nutrients requirements of growing rabbits according to NRC (1977) and Cheeke (1987). Diets were pelleted at investment unit, Poultry Production Department, Faculty of Agriculture, Mansoura University, Dakahlia Governorate, Egypt.

### **Digestibility trail and caecotrophy test.**

A total number of 84, unsexed, New Zealand White (NZW) weaned rabbits at 5 weeks old, were randomly divided into seven experimental groups (12 rabbits each) for growing trail lasted for 5-14 weeks of age. The data of first part of this study concerning the rabbits performance and blood parameters will be presented later one (Gabr *et al*, 2011). Seven digestion trails and caecotrophy tests were performed. At the end of the growing trail, a total number of 42 male rabbits were selected randomly from the previous herd and fed on the same dietary treatments (6 male for each treatment, 3 with collar to prevent caecotrophy and 3 without collar, to allow caecotrophy) and were allotted to meet the different treatments. The rabbits were housed individually in metabolic cages with a screen allowing facilitating the collecting faeces uncontaminated with urine through the digestibility trail. Quantitative collection of faeces was started 24 hrs after offering the daily feed. Faeces of each male were collected and feed intake was recorded daily in the morning for a collecting period 10 days was preceded by another 7 days as a preliminary period. The same feeding regimes used during the feeding trial (5-14weeks of age) were also followed through the digestibility trail. Collected



Duncan's multiple range test (Duncan, 1955) was also used for the comparison among means of the experimental groups.

## RESULTS AND DISCUSSION

### Chemical analysis of tested diets, dried onion (DO) and garlic (DG):

The determined and calculated chemical analysis of tested diets (Table 1&2) showed that all tested diets were practically similar, being formulated to be iso-nitrogenous and iso-caloric as designed. In addition, the chemical composition of DO and DG % was similar in DM, OM, CP and Ash contents, while EE in DO was higher than that of DG (0.68 vs 0.25%, respectively), and the reverse trend was true in case of CF (2.88 vs 5.27 %, respectively). Similar chemical analysis for DG have been recorded by Nwinuke *et al* (2005), Grela and Kelbaniuk (2007) and Otunola *et al* (2010) and for DO (Farral, 1985 and Nwinuka *et al* 2005).

**Table (1): Calculated composition of the basal diet.**

Ingredients:	Quantity (%)
Alfalfa hay	36
Yellow corn	10
Soybean meal(S.B.M)	8.6
Wheat bran	31
Barley grains	9.5
DL-methionine	0.1
Molasses	3.0
CaCo3	1.0
NaCl	0.5
Vitamin and mineral premix	0.3
Total	100
<b>Calculated diet composition:</b>	
Dry matter (D M ) %	91.09
Crude protein (%)	16.20
Ether extract (%)	2.79
Nitrogen free extract (NFE)%	49.9
Ash (%) <sup>2</sup>	8.10
DE (Kcal /Kg diet) <sup>5</sup>	2473.41
Crude fiber (%)	14.10
Neutral detergent fiber (NDF %)* <sup>3</sup>	38.19
Acid detergent fiber (ADF %)** <sup>3</sup>	22.29
Hemicellulose ( % ) <sup>4</sup>	15.9
Calcium (%) <sup>1</sup>	0.956
Phosphorus (%) <sup>1</sup>	0.527
Lysine (%) <sup>1</sup>	0.71
Sulfer amino acid (S .A.A%) <sup>1</sup>	0.60

1. Calculated according to Cheeke (1987)
2. Calculated according to NRC 1977,
3. Calculated according to Pagno Toscan *et al*, 1986 using the following equation:  
 \* NDF% = 28.924 + 0.657 (% crude fiber)      \*\*ADF%= 9.432 + 0.912 (%crude fiber)  
 \* NDF = cellulose +hemicellulose +lignin      \*\*ADF = cellulose + lignin
4. Hemicellulose (%) = NDF –ADF
5. Calculated according to Fekete (1987) using the following equation:  
 DE (kcal/kg)=[7.1(CP,g/kg)+12 ( EE,g/kg )+5.59 (NFE,g/kg )-1801]



**Dry matter intake (DMI), digestibility and nutritive values of tested diets:**

The obtained data (Table 3) indicated that dry matter intake ranged between 87.5 to 105.17g without significant differences among different dietary treatments. Also insignificant differences were detected between collared and uncollared rabbits. These results are in agreement with those of Ahmed *et al* (2005) and Dairo (2008).

Concerning nutrients digestibility coefficients, the highest significant ( $p<0.05$ ) value for DM was (74.77%) recorded for group receiving (0.5 % DO+ 0.5 %DG, while the lowest value was (66.93%) recorded with rabbits fed on 1.0% DO. Also, the highest CP digestibility coefficients were recorded with rabbits given 1.5 % DG, 1.5 % DO, 1 % DG and 0.5 % DO + 0.5 % DG diets, respectively, without significant differences among them, but they significantly higher ( $p<0.05$ ) than those of control, 1.0 % DO and 0.75% DO + 0.75 % DG. Digestibility coefficient of CF was significantly ( $p<0.05$ ) higher for rabbits fed 0.5 % DO + 0.5 % DG, than the other treated groups. The previous results are in agreement with those reported by El-Hindawy *et al* (2003) and Abdel-Azeem and Abdel-Reheem (2006).

As for the nutritive values of tested diets (Table 3), the highest TDN%, DE and ME Kcal/Kg diet were recorded with 0.5 % DO + 0.5 % DG which was significantly higher than those of 1.0 % DO and 1.5 % DO, but it was not significantly differed compared with control group, 1 % DG, 1.5 % DG, and 0.75% DO + 0.75 % DG. Regarding with DCP %, the highest DCP% values were recorded with 1.5 % DG, 1.5 % DO, 0.5 % DO + 0.5 % DG and 1 % DG without significant differences among them, but they were significantly ( $p<0.05$ ) higher than those of 1.0 % DO, control and 0.75% DO + 0.75 % DG without significant differences among the latter ones. The improvements effect of DO or/and DG supplementation on nutrients digestibility and hence the nutritive values of tested diets may be due to the presence of natural substances in the DO or/and DG which enhanced the activity of enzymes responsible for the digestion of nutrients in the digestive tract (El-Hindawy *et al*, 2003) as well as increase beneficial microbial activity. These findings may suggest that these supplements render the feeds more available for utilization, either by affecting positively the population of microflora or improving feed utilization through slowing feed rate of passage through the digestive tract, which was reflected later in better absorption (El-lamei, 2001).

**Coprophagy status effects:**

All nutrients digestibility coefficients as well as nutritive values of tested diets are shown in (Table 3), for rabbits allowed for coprophagy practice (uncollared) showed similar digestibility and nutritive values like those prevented coprophagy. Similar trend was observed by Mekawy (2007).

**Caecum pH and microflora:**

Results in Table (4) showed that mean of pH values for rabbits received control diet was insignificantly higher than those of the other dietary treatments. pH values of caecum appendix content are within the normal ranges reported by Lebas *et al* (1997). Similar trend have been reported by Abdel-Azeem and Abdel-Reheem (2006).



**Table (4): Least square means  $\pm$ SE of some factors affecting Caecum activity of rabbits for the experimental diets.**

Treatments	Caecum activity			
	pH	Bacterial total count**	E.coli total count	Salmonella and shigella
T1 Control diet	6.13 $\pm$ 0.040	23.50 $\pm$ 6.87 <sup>e</sup>	ND	ND
T2 (1% DO)	5.94 $\pm$ 0.040	27.50 $\pm$ 6.87 <sup>ed</sup>	ND	ND
T3(1.5% DO)	5.85 $\pm$ 0.040	172 $\pm$ 6.87 <sup>b</sup>	ND	ND
T4(1% DG)	5.89 $\pm$ 0.040	300.25 $\pm$ 6.87 <sup>a</sup>	ND	ND
T5(1.5% DG)	5.95 $\pm$ 0.040	47.50 $\pm$ 6.87 <sup>d</sup>	ND	ND
T6(0.5% DO+0.5 %DG)	5.93 $\pm$ 0.040	85.00 $\pm$ 6.87 <sup>c</sup>	ND	ND
T7(0.75% DO+ 0.75% DG)	5.89 $\pm$ 0.040	32.50 $\pm$ 6.87 <sup>ed</sup>	ND	ND

a, b, c, d, e, f and g Means with the same letter within each column are not significantly different.

SE=Standard error

ND=not detected

\*\* No. of bacterial cells / gm of caecum content (total count  $\times 10^5$ )

Rabbits fed diet containing 1.0 % DG recorded the highest means ( $p < 0.05$ ) of total bacterial count of caecum appendix contents followed by those given 1.5 % DO or 0.5 % DO + 0.5 % DG. The lowest means of total bacterial count were recorded by rabbits received control diet. The *E.coli* and *Salmonella* and *Shigella* counts seem to be undetected in all dietary treatments including control group (Table, 4). This may be due to the inclusion of the tested additives (garlic or onion) in rabbit diets, which contain active compounds such as allicin, quercetin, aflavonoid (one category of antioxidant compounds), (Abou EL-Wafa *et al.*, 2002). As well as they are rich in fructo-oligosaccharides which can selectively influence the intestinal microflora by either encourage the growth of beneficial bacteria and inhibit pathogenic species. The absent of the *E.coli* and *Salmonella* and *Shigella* bacteria in control group could be hardly explained on the light of the obtained results with other dietary treatments.

### Conclusion

It could be concluded that the tested botanical feed additive (dried onion or / and dried garlic) had a positive influence on most of digestibility coefficients and nutritive values of tested diets as well as showed healthy caecum activity of NZ W rabbits. The best level was mix of dried onion and dried garlic at 0.5% for each followed by 1.5% dried onion and 1.5% dried garlic.

## REFERENCES

- Abdel-Azeem, F. and Abdel-Reheem, M.A.T. (2006). Effect of supplemental copper and fresh garlic on performance and some biochemical changes in growing rabbits. *Egyptian J. of rabbit Sci.* 16(2): 341-366.
- Aboul-fotouh, G.E.; Allam, S.M.; Shehata, E.I. and Abdel-Azeem, S.N. (2000). Effect of some medicinal plants as feed additives on milk production and composition of lactating buffaloes. *Egyptian J. Nutrition and Feeds*, 3 (1): 31-41.



- Abou EL-Wafa, S.A.; Sedki, A.A. and Ismail, A.M. (2002). Response of growing rabbits to diet containing black seed, garlic or onion as natural feed additives. *Egyptian J of Rabbit-Sci*, 12(1):69-83.
- Ahmed, A. A.; Bassuony, Neamat I.; Awad, Set El-Habiab, S.; Aiad, A.M. and Mohamed, S.A. (2009). Adding natural juice of vegetables and fruitage to ruminant diets (B) nutrients utilization, microbial safety and immunity, effect of diets supplemented with lemon, onion and garlic juice fed to growing buffalo calves. *World J. of Agric. Sci.*, 5(4):456-465.
- Ahmed, F.G.; Yacout, M.H. and Abo-Donia, F.M. (2005). Effect of using eucalyptus globules leaves in growing rabbit diets. *Egyptian J. of Rabbit Sci.*, 15(1):1-11.
- Aiad, A.M.; Bassuony, N.I.; Afifi, A.A. and Abo-Donia, F.M. (2008). Adding natural juice of vegetables and fruitage to ruminant diets: (A) lemon, onion and garlic juice supplement to diets fed to suckling Buffalo calves and its effect on digestibility, growth performance and fungi count. *World J. of Agric. Sci.* 4(2): 149-156.
- A.O.A.C. (2000). Association of Official Analytical Chemists. W. Horwitz (Editor) official methods of analysis, 13<sup>th</sup>, Washington, D.C.
- British Standard Institution (1991). Microbiological examination of food and animal feeding stuff. Part I.
- Cheeke, P.R. (1987). Rabbit feeding and nutrition. Academic Press. Orln do, Florida, U.S.A.
- Cheeke, P.R.; Patton, N.M. and Tempelton, G.S. (1982). Rabbit production 5<sup>th</sup> Ed. Int. Print. and Publ. Danville Il.
- Dairo, F.A.S. (2008). Assessment of loofah gourd seeds *luffa cylindrical* (Roem) on performance and some haematological indices of rabbit weaners. 9<sup>th</sup> World Rabbit Congress-June5 10-13, 2008- Verona-Italy.
- Difco laboratories Incorporated (1984). Difco manual of dehydrated culture media and reagents for microbiological and clinical laboratory procedure, 10<sup>th</sup> Ed. Michigan, USA.
- Duncan, D.B. (1955). Multiple ranges and multiple F-test Biometrics, 11: 1-42.
- El-Ashry, M.A.; El-Bordeny, N.E.; Khattab, H.M. and El-Sayed, H.M (2006). Effect of diet supplemented with medicinal herbs on nutrient digestibility and some blood metabolites of buffalo calves. *Egyptian J. Nutrition and Feeds*, 2: 179-191.
- Ellelamei, R.R. E. (2001). Evaluation of some medical plants in rabbit nutrition. M.Sc thesis, Fac. of Agric., Cairo Univ.
- El-Hindawy, M.M.; Yacout, M.H.; Hassa, I.I. and ebidalla, N.A. (2003). Evaluation of onion seeds as unconventional feed stuff for growing rabbit. *Egyptian J. of Rabbit Sci.* 13(2): 183-194.
- Farral, K.T. (1985). Spices and condiments and seasoning. AVI Publishing, West Port, CT.
- Fekete, S. (1987). The new Hungrain system for evaluation of feed energy. Proceedings of 1<sup>st</sup> North American Rabbit Congress. October, 10-13, Portland-OR, USA.

- Gabr,A.A.; El-Sayaad, G.A.; Abdel-Azeem, A.F and Ibrahim, W.A.(2011). Productive performance,some blood constituents and immunological response of Rabbit fed diets supplemented with different levels of onion and garlic as safe feed additives. Egyptian J. of Nutrition and Feeds. The 13<sup>rd</sup> conference of animal feeding, 22-25 November, Marsa Alam, Egypt.(Under publication).
- Grela,E.R. and Klebaniuk R. (2007). Chemical composition of garlic preparation and its utilization in piglet diets. *Medycyna Wet.*, 63(7): 792-795.
- Gupta,N.; Kumar,A. and Tiwar, D.P. (2005). Effect of herbs as feed additives on nutrient utilization and growth in crossbred heifers fed paddy straw ration. *Indian J. of Animal Sci.*,75(1):52-55.
- Helal, F.I.S. and Mohamed, R.S. (2001). Influence of garlic oil (*Allium Sativum* L.) and some of sulfur containing constituents on some biochemical changes in milk, blood and feed utilization in rabbits, with especial reference to its hypocholesterolemic activity. *Egyptian J. Nutrition and Feeds*, 4:909-918.
- Kumar,M. and Berwal, J.S. (1999). Sensivity of food pathogens to garlic (*Allium sativum*). *3 of Appl. Microbiol.*, 84: 213-215.
- Lebas, F.: Coudert, P.: Rochambean, H. and Thebauff, R. G. (1997). Nutrition and feeding in the rabbit. *Husbandry, Health and production*, pp 19-21.
- Line-Eric, J.; Stan Bailey, J.; Nelson, C.A.; Norman, S.J. and Thomas, T. (1998).Effect of supplemented feed on *Salmonella* and company lobacter population in broilers. *Poultry Sci.*, 77:405.
- Mekawy, H.M.S.H. (2007). The possibility of using irradiated untraditional feed-stuffs in rabbits nutrition. Ph.D. Thesis, Fac. Of Agric., Moshtohor, Benha univ., Egypt.
- National Research Council (NRC, 1977). Nutrient requirements of domestic animals. Nutrient Requirements of Rabbits. USA National Academy of Science. Washington, D.C.
- Nwinuka, N. M; Ibeh, G. O. and Ekeke, G. I. (2005). Proximate composition and levels of some toxicants in four commonly consumed spices. *J. Appl. Sci. Environ. Mgt*, 9(1): 150-155.  
(C.F. Textetbook of rabbit medicine, Elsevier Science Limited 2002)
- Otunola, G.A.; Oloyede, O.B.; Oladiji, A.T. and Afolayan, A.J. (2010). Compative analysis of the chemical composition of three spices-*Allium Sativum* L. *zingiber officinale* Rosc. and *capsicum frutescens*l. commonly consumed in Nigeria. *Afr. J. of Biotechnology*, 9 (41):6927-6931.
- Pagano Toscano, G.; Benatti, G. and Zoccarato, I. (1986). Comparison of crude fiber and the Van Soest detergent methods for fiber determination in rabbit feeds. *J. Appl. Rabbit Res.* 9: 69-75.
- Pond, W.G.; Church, D.C.; Pond, K.R..and Schoknecht, P.A. (2006). Basic animal nutrition and feeding. Fifth Edition. Nice printing press, Delhi.
- Saleem, Z. M. and Al-Delaimy, K.S. (1982).Inhibition of *Bacillus aurous* by garlic extracts. *J. of food protection*, 45:1007-1009.
- SAS (2002). SAS user's guide Ins. Inc. Cary. NC. USA.

- Schiemann, R.; Nehring, L.; Hoffmann, W.; j. and Chudy A. (1972). Energetische Futterbewertung and Energianormen, VEB Deutscher Landwirtschaftsverlag, Berlin. P.72
- Wager, A.; Ouaratulain, S.; Altaf, H.; Ahmed, G.M. and Ashhar, Z. (1994). Evaluation of different garlic extracts for antibacterial activity. Sci. Intl., 5:385-386.

**تأثير اضافة مستويات مختلفه من البصل والثوم المجفف ومخاليطهم على معاملات الهضم وظاهرة الاجترار الكاذب Coprophagy ونشاط الاعور في الأرانب**  
**جمال على الدين الصياد<sup>1</sup> ، احمد عبد الرازق جبر<sup>2</sup> ووحيد عطيه الطيب ابراهيم<sup>2</sup>**  
**1- كلية الزراعة - بمشهر جامعه بنها**  
**2- كلية الزراعة - جامعه المنصوره**

اجريت هذه الدراسه بمحطه تجارب قسم الانتاج الحيوانى - كلية الزراعة بمشهر - جامعه بنها وقد استخدمت فى هذه الدراسه عدد 42 ذكر نيوزلندى تم توزيعها عشوائيا الى 7 مجموعات 6 ذكر (3 بطوق و 3 بدون طوق) كالتالى : 1- المجموعه الاولى تغذت على العليقه الكنترول 2- المجموعه لثانيه تغذت على العليقه الكنترول مضافه اليها 1% من البصل المجفف ، 3- المجموعه الثالثه تغذت على العليقه الكنترول مضاف اليها 1.5 بصل مجفف ، 4- المجموعه الرابعه ، تغذت على العليقه الكنترول مضاف اليها 1% ثوم مجفف ، 5- المجموعه الخامسه ، تغذت على العليقه الكنترول مضاف اليها 1.5% ثوم مجفف ، 6- المجموعه السادسه ، تغذت على العليقه الكنترول مضاف اليها 0.5% بصل مجفف + 0.5% ثوم مجفف ، 7- المجموعه السابعه ، تغذت على العليقه الكنترول مضاف اليها 0.75% بصل مجفف + 0.75% ثوم مجفف.

**وتتلخص أهم النتائج فى هذه التجربه:**

**المأكول من الماده الجافه:**

- 1- أشارت النتائج المتحصل عليها إلى عدم وجود فروق معنويه فى المأكول من الماده الجافه بين مجموعه الكنترول مقارنة بالمعاملات الاخرى.
- 2 - أظهرت النتائج عدم وجود فرق معنوى فى المأكول من الماده الجافه بين مجموعه الأرانب التى منعت من ظاهرة الاجترار الكاذب Coprophagy مقارنة بالتى لم تمنع.

**معاملات الهضم:**

- 1- اتضح من النتائج المتحصل عليها عدم وجود فرق معنوى بين معاملات هضم كل من الماده العضويه OM والمستخلص الأثيرى EE والكربوهيدرات الذائبه NFE عندما تناولت الأرانب العليقه الكنترول مقارنة بتلك التى تغذت على العلائق الاخرى.
- 2- اظهرت النتائج وجود فرق معنوى لمعامل هضم البروتين بمجموعات الأرانب المغذاه على العليقه الكنترول مضافا " اليها 1.5% ثوم و 1.5% بصل ومخلوط من البصل والثوم بمستوى 0.5% مقارنة بالكنترول والمجموعات الاخرى.
- 3- أدى اضافة مخلوط من البصل والثوم عند مستوى 0.5% لكل منهما فى علائق الأرانب الى تحسن معنوى لمعامل هضم الالياف CF .
- 4- اشارت النتائج المتحصل عليها الى تحسن معنوى عند مستوى معنويه 5% فى القيم الغذائيه (مجموع المركبات الغذائيه المهضومه TDN والطاقه المهضومه DE والطاقه الممتله ME عندما تناولت الأرانب عليقه الكنترول مضافا " اليها مخلوط البصل والثوم عند مستوى 0.5% لكل منهما عند مقارنتها بتلك التى تغذت على 1% بصل مجفف.
- 5- اوضحت النتائج المتحصل عليها عدم وجود فروق معنويه فى معاملات الهضم المختلفه ( الماده الجافه DM والماده العضويه OM والبروتين الخام CP والدهون EE والالياف الخام CF والكربوهيدرات الذائبه NFE ) والقيم الغذائيه (مجموع المركبات الغذائيه

المهضومه TDN والطاقة المهضومه DE والطاقة الممثله ME ) عند مقارنه مجموعه الأرانب التي منعت من ظاهرة الاجترار الكاذب Coprophagy بتلك التي سمح لها بالاجترار الكاذب (تناول الروث الطرى).

**نشاط الأعر:**

أدى ادخال الاضافات فى علائق الأرانب الى زيادة العدد الكلى من بكتريا الاعور مقارنه بالكنترول مع غياب البكتريا المرضيه ( الايشريشيا كولاي والسالمونيلا والشيجلا) فى محتويات الاعور. وعموما نستخلص من هذه الدراسه انه يمكن التوصيه بإضافه مسحوق الثوم و البصل إلى علائق الأرانب لتحسين الكفاءة الهضمية وصحة وسلامة الجهاز الهضمى ونشاط الأعر، وقد كانت أفضل المستويات المستخدمة المخلوط المكون من البصل والثوم عند مستوى 0.5 % لكل منهما يليه المستوى 1.5 % بصل و 1.5 % ثوم.

**قام بتحكيم البحث**

كلية الزراعة – جامعة المنصورة  
كلية الزراعة – جامعة الأزهر

أ.د / محمود يوسف العايق  
أ.د / رضا سلامه محمد



Table (3): Least square means and standard errors for digestibility coefficients and nutritive values of the tested diets.

Items	Control diet (T1)	Levels dietary treatment						Coprophagy status		Standard errors		Significant levels	
		Level of dried onion (DO)		Level of dried garlic (DG)		Mixture of (DO and DG)		Collared Rabbits	Un collared Rabbits	treatments	coprophagy	treatment	coprophagy
		T2 (1%)	T3 (1.5%)	T4 (1%)	T5 (1.5%)	T6 (0.5%+ 0.5%)	T7 (0.75%+ 0.75)						
No. of rabbits	6	6	6	6	6	6	6	21	21				
Dry matter intake(DMI) g/h/d	105.17	100.83	91.33	87.5	101.83	88.67	92.83	94.96	96.14	4.99	2.67	NS	NS
<b>Digestibility coefficients %</b>													
DM	70.64 <sup>abc</sup>	66.93 <sup>c</sup>	68.81 <sup>bc</sup>	72.49 <sup>ab</sup>	71.63 <sup>ab</sup>	74.77 <sup>a</sup>	69.47 <sup>bc</sup>	71.27	69.93	1.53	0.82	*	NS
OM	71.36	68.19	69.73	73.13	71.66	74.96	71.02	72.04	70.83	1.45	0.77	NS	NS
CP	68.19 <sup>b</sup>	68.82 <sup>b</sup>	79.54 <sup>a</sup>	78.60 <sup>a</sup>	80.11 <sup>a</sup>	79.02 <sup>a</sup>	72.54 <sup>b</sup>	74.83	75.54	1.85	0.99	**	NS
CF	52.01 <sup>bcd</sup>	42.97 <sup>d</sup>	46.87 <sup>cd</sup>	56.52 <sup>b</sup>	56.22 <sup>bc</sup>	60.80 <sup>a</sup>	54.25 <sup>bc</sup>	51.41	55.64	2.98	1.59	**	NS
EE	76.81	74.36	78.69	71.29	75.96	71.43	71.09	74.22	74.23	2.76	1.47	NS	NS
NFE	77.89	75.62	72.82	76.27	73.13	76.46	75.54	75.9	74.87	1.51	0.81	NS	NS
<b>Nutritive values%</b>													
TDN <sup>1</sup>	66.92 <sup>ab</sup>	62.92 <sup>b</sup>	64.18 <sup>b</sup>	67.29 <sup>ab</sup>	66.98 <sup>ab</sup>	69.51 <sup>a</sup>	66.13 <sup>ab</sup>	65.73	66.19	1.35	0.72	*	NS
DCP	12.44 <sup>b</sup>	12.38 <sup>b</sup>	14.47 <sup>a</sup>	14.31 <sup>a</sup>	14.58 <sup>a</sup>	14.38 <sup>a</sup>	13.20 <sup>b</sup>	13.61	13.74	0.34	0.18	**	NS
DE <sup>2</sup>	2946.67 <sup>abc</sup>	2777.33 <sup>c</sup>	2853.17 <sup>bc</sup>	2981.00 <sup>ab</sup>	2972.17 <sup>ab</sup>	3076.00 <sup>a</sup>	2921.17 <sup>abc</sup>	2910.33	2954.66	59.08	31.58	*	NS
ME <sup>3</sup>	2828.66 <sup>abc</sup>	2666.17 <sup>c</sup>	2739.00 <sup>bc</sup>	2861.50 <sup>ab</sup>	2853.00 <sup>ab</sup>	2953.17 <sup>a</sup>	2804.33 <sup>abc</sup>	2793.9	2836.33	56.68	30.29	*	NS

a ,b and c means within each row with different superscripts are significantly ( p<0.05) different. NS= not significant

1-Total digestible nutrient (TDN) was calculated according to the classic formula of (Cheeke *et al*, 1982) as follows:TDN%=%DCP+%DCF+%DNFE+%DEE x 2.25

2-Digestible energy (DE) was calculated according to Schiemann *et al.*, (1972) as follows: DE (Kcal/kg diet) =5.28 (DCP, g kg) + 9.51(DEE, g/kg) +4.20 (DCF, g/kg) + 4.20 (DNFE, g/kg)

3-Metabolizable energy (ME) was calculated according to (Pond *et al*, 2006) as follows: ME (in kcal/kg) =DE (in kcal/kg) x 0.96-(0.202 x protein %) / 100

**Table (2): Chemical analysis of tested diets and natural feed additives (dried garlic DG and dried onion DO powder on DM basis %).**

Items	Control diet ( T1 )	Levels dietary treatments						Chemical analysis of dried garlic (DG) and dried onion (DO) powder	
		Levels of dried onion(DO)		Levels of dried garlic(DG)		Mixture of (DO) and (DG)		Dried garlic powder	Dried onion powder
		T2 (1%)	T3 (1.5%)	T4 (1%)	T5 (1.5%)	T6 (0.5%+0.5%)	T7 (0.75%+0.75%)		
Dry matter (DM %)	91.09	91.12	91.16	90.96	91.20	91.15	91.06	92.76	92.40
Crude protein(CP % )	18.19	18.21	18.24	18.13	18.25	18.12	18.18	14.96	13.79
Ether extract (EE %)	3.13	2.89	2.85	2.93	2.88	2.79	2.95	0.25	0.68
Ash ( % )	10.52	10.67	10.59	10.68	10.40	10.34	10.58	3.50	3.70
Organic matter (OM %)	89.48	89.33	89.41	89.32	89.60	89.66	89.42	96.50	96.30
Nitrogen free extract (NFE )	52.38	52.42	52.56	52.47	52.73	52.95	52.54	76.02	78.95
DE(Kcal /Kg diet)	2794.13	2768.99	2774.14	2770.90	2787.96	2780.23	2780.77	3540.68	3673.00
Crude fiber (CF %)	15.78	15.81	15.76	15.79	15.74	15.80	15.75	5.27	2.88
Neutral detergent fiber(NDF %)	39.29	39.31	39.28	39.30	39.27	39.30	39.27	32.39	30.82
Acid detergent fiber(ADF%)	23.82	23.85	23.81	23.83	23.79	23.84	23.80	14.24	12.06
Hemicellulose ( % )	15.47	15.46	15.47	15.47	15.48	15.46	15.48	18.15	18.76